

Conflicts of interest

Dr. Barroso reports personal fees from AstraZeneca, Boehringer Ingelheim, Lda, Eli Lilly and Company, Pierre Fabre Portugal, Roche Farmacêutica Química, Lda, Merck and Bristol-Myers Squibb, outside the submitted work. The other authors have no conflicts of interest to declare.

Acknowledgements

None.

References

1. Dirican N, Dirican A, Anar C, Atalay S, Ozturk O, Bircan A, et al. A new inflammatory prognostic index based on C-reactive protein, the neutrophil to lymphocyte ratio and serum albumin is useful for predicting prognosis in non-small cell lung cancer cases. *Asian Pac J Cancer Prev.* 2016;17:5101–6.
 2. Shrotriya S, Walsh D, Bennani-Baiti N, Thomas S, Lorton C. C-reactive protein is an important biomarker for prognosis tumor recurrence and treatment response in adult solid tumors: a systematic review. *PLoS ONE.* 2015;10:e0143080.
 3. Derman BA, Macklis JN, Azeem MS, Sayidine S, Basu S, Batus M, et al. Relationships between longitudinal neutrophil to lymphocyte ratios, body weight changes, and overall survival in patients with non-small cell lung cancer. *BMC Cancer.* 2017;17:141.
 4. Forget P, Khalifa C, Defour JP, Latinne D, Van Pel MC, De Kock M. What is the normal value of the neutrophil-to-lymphocyte ratio? *BMC Res Notes.* 2017;10:12.
 5. Zhang F, Ying L, Jin J, Chen K, Zhang N, Wu J, et al. The C-reactive protein/albumin ratio predicts long-term outcomes of patients with operable non-small cell lung cancer. *Oncotarget.* 2017;8:8835–42.
 6. Gomes M, Teixeira AL, Coelho A, Araújo A, Medeiros R. The role of inflammation in lung cancer. *Adv Exp Med Biol.* 2014;816: 1–23.
 7. Jing X, Huang C, Zhou H, Li C, Fan L, Chen J, et al. Association between serum C-reactive protein value and prognosis of patients with non-small cell lung cancer: a meta-analysis. *Int J Clin Exp Med.* 2015;8:10633–9.
 - D. Machado ^{a,*}, C. Marques ^a, M. Dias ^b, S. Campainha ^b, A. Barroso ^b
- ^a Pulmonology Department, Gaia/Espinho Hospital Center, Vila Nova de Gaia, Portugal
^b Thoracic Tumors Multidisciplinary Unit, Pulmonology Department, Gaia/Espinho Hospital Center, Vila Nova de Gaia, Portugal
- * Corresponding author.
E-mail address: daniela.pc.machado@gmail.com (D. Machado).
- <https://doi.org/10.1016/j.pulmoe.2019.02.007>
2531-0437/© 2019 Published by Elsevier España, S.L.U. on behalf of Sociedade Portuguesa de Pneumologia. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Immunotherapy in Allergic Asthma – 5 year analysis: Is it a curative approach?



Dear Editor,

Asthma is a common, chronic and heterogeneous disease; it affects people of all ages and there has been a recent increase in its prevalence and severity. It may be mild, barely noticed by the patient, or it may range all the way to very severe disease, causing constant symptoms that greatly affect the quality life of the patient. The long-term goals of asthma management are to achieve good symptom control and minimize future risk of exacerbation, fixed air flow limitation and treatment of side effects.¹ The patient own goals regarding their asthma and its treatment should also be identified. During the last decade several new drugs addressing asthma have been developed and approved.

The immune system is a complex interactive network with the capacity of protecting the host from a number of pathogens while maintaining a state of tolerance to self and innocuous exogenous antigens. Allergy is one of the immune tolerance-related diseases that arises as a direct consequence of a dysregulated immune response.

Currently, allergen-specific immunotherapy (allergen-SIT) remains the single curative approach to allergic diseases with the potential to modify its course.² It is proven to be

the only therapy that alters the natural history of allergic disease, prevents progression, prevents the development of new sensitizations and may even prevent the development of asthma in patients with allergic rhinitis.³ It consists of administering gradually increasing doses of a specific allergenic extract to an allergic subject (with specific documented IgE), at an effective dose, in order to decrease the symptoms associated with subsequent exposure to the causative allergen.³ Multiple mechanisms are involved in the suppression and/or control of allergic inflammation.⁴ The aim of allergen-SIT is to induce the peripheral T cell tolerance, modulate the thresholds for mast cell and basophil activation and decrease IgE-mediated histamine release.⁴ Peripheral T cell tolerance is characterized by the generation of allergen-specific T-regulatory cells (Treg) that are able to produce anti-inflammatory cytokines such as IL-10 and TGF-β.⁵ Treg cells not only diminish Th2 immune responses, but also target other cell types such as mast cells, basophils, eosinophils, allergen-specific-IgE and are capable of inducing IgG4 and IgA production.⁶

A review of Abramson et al., which included eighty-eight trials, showed a significant reduction in asthma symptoms, reduction in use of medications and improvement in bronchial hyper-reactivity following immunotherapy.⁷ According to the Global Initiative for Asthma (GINA) 2017, for patients with allergic rhinitis and sensitization to house dust mite, with exacerbations despite low-high dose of ICS, we must consider adding sublingual allergen immunotherapy

Table 1

	Year before the beginning of SIT (I)	1 year after the start of SIT (J)	3 years after the start of SIT (K)	5 years after the start of SIT (L)	p ^a	η^2 partial	Dif. I-Jp ^b	Dif. I-Kp ^b	Dif. I-Lp ^b	Dif. J-Kp ^b	Dif. J-Lp ^b	Dif. K-Lp ^b
No. exacerbations/year (Mean \pm SD)	3.90 \pm 2.72	0.14 \pm 0.36	0.05 \pm 0.22	0.19 \pm 0.40	<0.001	0.670	<0.001	<0.001	<0.001	0.909	0.999	0.709
CARAT (Mean \pm SD)	18.2 \pm 2.29	22.8 \pm 2.93	24.2 \pm 2.76	23.8 \pm 4.03	<0.001	0.513	<0.001	<0.001	<0.001	0.426	0.845	0.999
GINA therapeutic step (Median) (Min; Max)	3 (1; 4)	3 (1; 4)	3 (1; 4)	3 (1; 4)	0.282	-	-	-	-	-	-	-
FEV1% predicted (Mean \pm SD)	97.3 \pm 12.78	97.8 \pm 8.43	102.6 \pm 11.08	95.62 \pm 11.22	0.295	0.078	1.000	0.434	1.000	0.654	0.987	0.474
PEF% predicted (Mean \pm SD)	84.5 \pm 28.23	100.1 \pm 21.80	95.3 \pm 12.09	98.9 \pm 13.40	0.101	0.128	0.564	0.361	0.332	1.000	1.000	1.000

^a ANOVA of repeated measurements, MANOVA (used when the assumption of sphericity was not assumed and epsilon of Huynh–Feldt <0.7) or Friedman's test.

^b Sidak test.

(SLIT), provided Fev1 is >70% predicted.¹ Thus, in the latest GINA recommendations, allergen-SIT have been included for patients with exacerbation despite taking step 3 or step 4 therapy, as an add-on therapy.¹

Here, we do a prospective analysis of patients with asthma submitted to allergen-SIT followed in allergology appointment during 5 years (2013–2017). Statistical analysis was performed using the IBM SPSS® statistical program, version 25. The variables were described with mean, median, standard deviation, maximum and minimum. All the tests were considered significant when *p*-value did not exceed 0.05.

We included 35 patients with mean age 28.1 ± 11.4 years. The majority (94.3%) had associated allergic rhinitis. Regarding therapy 29% of the patients performed allergen-SIT for dermatophagoides pteronyssinus (Dpt), 23% Dpt/dermatophagoides farinae, 14% Dpt/Lepidoglyphus destructor and 31% for Gramineae mixture. The majority had allergenic polysensitization (57%).

In the year before the start of allergen-SIT, patients presented an average of 3.9 ± 2.7 exacerbations/year and mean CARAT of 18 ± 2.3 . One year after the onset of allergen-SIT, they presented a mean of 0.14 ± 0.36 exacerbations/year and mean CARAT of 22.8 ± 2.9 . At 3 years, they presented a mean of 0.05 ± 0.22 exacerbations/year and mean CARAT of 24.2 ± 2.7 , whereas at 5 years, they presented a mean of 0.19 ± 0.40 exacerbations/year and mean CARAT of 23.8 ± 4 . During this period they were performing therapy on step 3 of the GINA. There were statistically significant differences in the number of exacerbations/year and the CARAT questionnaire mean at 1, 3 and 5 years (*p* < 0.001). There were no statistically significant differences in bronchodilator and anti-inflammatory therapy during the same period (*p* = 0.282), Table 1. Regarding safety, most local adverse reactions were mild and there were neither life-threatening systemic reactions nor fatal events.

In conclusion, after one year of allergen-SIT, we verified a significant symptomatic improvement in the clinical control of asthma, as evidenced by the statistically significant difference in the number of exacerbations/year and in the CARAT questionnaire. This control remains stable after 3 and 5 years of therapy, without significant differences in bronchodilator and anti-inflammatory therapy. Therefore, the current evidence provides support for the efficacy and safety of allergen-SIT in the treatment of respiratory allergy, namely in clinical control of allergic asthma and rhinitis, representing a safe alternative or an add-on therapy to conventional inhaler therapy. However, it is essential to select

patients to undergo this treatment and which extracts to be used to optimize the cost/benefit.

Conflicts of interest

The authors have no conflicts of interest to declare.

References

1. Global Initiative for Asthma. Global strategy for asthma management and prevention; 2017. <http://ginasthma.org/2017-gina-report-global-strategy-for-asthma-management-and-prevention>
2. Muraro A, Roberts G, Agache I, Angier L, Bonertz A, Dhami S, et al. Allergen immunotherapy guidelines. European Academy of Allergy and Clinical Immunology (EAACI); 2017.
3. Nelson H, et al. Some highlights of the first century of immunotherapy, vol. 107. 2011 American College of Allergy; 2011.
4. Akdis C, Akdis M. Mechanisms of allergen-specific immunotherapy. *J Allergy Clin Immunol*. 2011;127:18–27, quiz 28–9.
5. Kearley J, Barker J, Robinson D, Lloyd C. Resolution of airway inflammation and hyperreactivity after *in vivo* transfer of CD4+CD25+ regulatory T cells is interleukin 10 dependent. *J Exp Med*. 2005;202(11):1539–47.
6. Kearley J, Robinson D, Lloyd C. CD4+CD25+ regulatory T cells reverse established allergic airway inflammation and prevent airway remodeling. *J Allergy Clin Immunol*. 2008;122:617–24.
7. Abramson M, Puy R, Weiner J. Injection allergen immunotherapy for Asthma. *Cochrane Database of Systematic Reviews*. 2010;(Issue 8.). Art. No.: CD001186.

J. Coutinho Costa^{a,*}, J.N. Machado^a, C. Ferreira^a, J.M.R. Gama^b, T. Almeida^a, A.M. Arrobas^a

^a *Pneumology B Unit, University Hospital Centre of Coimbra – Hospital Geral, Coimbra, Portugal*

^b *Centre of Mathematics and Applications, University of Beira Interior, Covilhã, Portugal*

* Corresponding author.

E-mail address: josecoutinhocosta99@gmail.com (J. Coutinho Costa).

<https://doi.org/10.1016/j.pulmoe.2019.01.007>

2531-0437/

© 2019 Sociedade Portuguesa de Pneumologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).