

Covid-19: Round and oval areas of ground-glass opacity



The World Health Organization has classified the COVID-19 situation as a pandemic. Patients infected with COVID-19 typically present with fever, cough, dyspnea, and myalgia, and the infection may cause severe pneumonia. Laboratory findings are unspecific. Although test results are normal for many patients, the predominant laboratory abnormalities include the elevation of inflammatory markers, such as C-reactive protein, lactate dehydrogenase, and the erythrocyte sedimentation rate. Additionally, lymphopenia is consistently present in more than 40% of patients.¹⁻³ Currently, real-time reverse-transcription polymerase chain reaction is the reference standard test for the definitive diagnosis of COVID-19 infection.

A 44-year-old man presented to the hospital (Centro Hospitalar do Porto, Porto, Portugal) on 7 March 2020 with a 6-day history of fever, cough, rhinorrhea, diffuse myalgia and fatigue. He had had close contact with a friend who had traveled to Milan, Italy, on 22–23 February 2020, during the COVID-19 pneumonia outbreak. On admission, the patient was in good general condition; he was tachypneic (respiratory rate of 30 breaths/min), his body temperature was 38.2 °C, and cardiac auscultation was normal, with no murmur or arrhythmia. Pulmonary auscultation demonstrated the presence of sparse bilateral crackles. Laboratory tests showed a normal blood cell count, erythrocyte sedimentation rate of 84 mm/h (normal = 0–10 mm/h), C-reactive

protein level of 14 mg/L (normal = 0.3–10 mg/L), and unremarkable lactate dehydrogenase, creatine phosphokinase, and liver function findings. Blood gas analysis yielded normal findings (O_2 saturation = 98%). The patient's respiratory rate returned to normal after the normalization of his temperature. Non-enhanced chest computed tomography (CT) showed multiple round and oval ground-glass opacities in both lungs, with a crazy-paving pattern (Fig. 1B–D). No mediastinal lymphadenopathy or pleural effusion was present. Real-time reverse-transcription polymerase chain reaction of a nasopharyngeal sample revealed positivity for 2019-nCoV nucleic acid. Hydroxychloroquine and symptomatic medication were administered. The patient recovered uneventfully and was discharged after 16 days in an asymptomatic state.

Although our patient had pneumonia, as confirmed by CT, the chest radiograph was normal. Chest radiography has not been recommended as a first-line imaging modality for the diagnosis of COVID-19 due to its limited sensitivity in the detection of ground-glass opacities and other incipient pulmonary findings of the infection, which are evident on CT. However, nonspecific chest radiography findings have been reported occasionally, particularly for patients with severe disease.^{4,5} The role of CT in COVID-19 evaluation is the subject of much discussion. Some authors suggest that CT has a pivotal role, whereas other investigators are less optimistic. The predominant CT findings are multifocal, bilateral, peripheral, and basal-predominant ground-glass opacities, often with round and/or oval morphology and/or consolidation. The crazy-paving pattern may be observed.




Figure 1 (A) A posteroanterior chest radiograph was considered normal. Unenhanced chest computed tomography with axial (B), coronal (C) and sagittal (D and E) maximum-intensity projection imaging demonstrated areas of ground glass opacity, many with round and oval morphologies, in both lungs. Not also in B inter- and intralobular septal thickening with a crazy-paving pattern (arrows).

This pattern is defined as thickened interlobular septa and intralobular lines superimposed on a background of ground-glass opacities. Pleural effusion, small lung nodules, cavitation, and lymphadenopathy are very uncommon findings.^{2,5,6} These CT findings are not specific to COVID-19; similar results can be obtained for other infectious and non-infectious diseases.⁶

However, two characteristics of the ground-glass opacities may suggest the diagnosis of COVID-19 in the context of the current pandemic. The presence of multifocal nodular (round or oval) ground-glass opacities^{7,8} and/or the association of these opacities with reticulation (the crazy-paving pattern)^{8–10} should alert the radiologist to the possibility of COVID-19 infection. The latter finding appears particularly when the disease progresses. Our patient presented both findings. We believe that these two findings are important for the diagnosis of COVID-19, although the crazy-paving pattern is less specific; Amorim et al.¹¹ observed it in 15% of 70 patients with confirmed H1N1 infection.

Conflicts of interest

The authors have no conflicts of interest to declare.

References

- Rodriguez-Morales AJ, Cardona-Ospina JA, Gutiérrez-Ocampo E, Villamizar-Peña R, Holguin-Rivera Y, Escalera-Antezana JP, et al. Clinical, laboratory and imaging features of COVID-19: a systematic review and meta-analysis. *Travel Med Infect Dis.* 2020;101623, <http://dx.doi.org/10.1016/j.tmaid.2020.101623>.
- Xiong Y, Sun D, Liu Y, Fan Y, Zhao L, Li X, Zhu W. Clinical and high-resolution CT features of the COVID-19 infection: comparison of the initial and follow-up changes. *Invest Radiol.* 2020, <http://dx.doi.org/10.1097/RLI.0000000000000674>.
- Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med.* 2020, <http://dx.doi.org/10.1056/NEJMoa2002032>.
- Zu ZY, Jiang MD, Xu PP, Chen W, Ni QQ, Lu GM, et al. Coronavirus disease 2019 (COVID-19): a perspective from China. *Radiology.* 2020;200490, <http://dx.doi.org/10.1148/radiol.2020200490>.
- Ng M-Y, Lee EY, Yang J, Yang F, Li X, Wang H, et al. Imaging profile of the COVID-19 infection: radiologic findings and literature review. *Radiol Cardiothorac Imaging.* 2020;2:e200034, <http://dx.doi.org/10.1148/ryct.2020200034>.
- Hope MD, Raptis CA, Shah A, Hammer MM, Henry TS, on behalf of six signatories. A role for CT in COVID-19? What data really tell us so far. *Lancet.* 2020;26, [http://dx.doi.org/10.1016/S0140-6736\(20\)30728-5](http://dx.doi.org/10.1016/S0140-6736(20)30728-5).
- Kong W, Agarwal PP. Chest imaging appearance of COVID-19 infection. *Radiol Cardiothorac Imaging.* 2020;2, <http://dx.doi.org/10.1148/ryct.2020200028>.
- Chung M, Bernheim A, Mei X, Zhang N, Huang M, Zeng X, et al. CT imaging features of 2019 novel coronavirus (2019-nCoV). *Radiology.* 2020;295:202–7, <http://dx.doi.org/10.1148/radiol.2020200230>.
- Pan F, Ye T, Sun P, et al. Time course of lung changes on chest CT during recovery from 2019 novel coronavirus (COVID-19) pneumonia. *Radiology.* 2020;200370, <http://dx.doi.org/10.1148/radiol.2020200370>.
- Zhou S, Wang Y, Zhu T, Xia L. CT features of coronavirus disease 2019 (COVID-19) pneumonia in 62 patients in Wuhan, China. *AJR Am J Roentgenol.* 2020;1–8, <http://dx.doi.org/10.2214/AJR.20.22975>.
- Amorim VB, Rodrigues RS, Barreto MM, Zanetti G, Hochhegger B, Marchiori E. Influenza A (H1N1) pneumonia: HRCT findings. *J Bras Pneumol.* 2013;39:323–9, <http://dx.doi.org/10.1590/S1806-37132013000300009>.

W. Schmitt^a, E. Marchiori^{b,*}

^a *Imaging Department - Centro Hospitalar do Porto, Porto, Portugal*

^b *Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil*

* Corresponding author.

E-mail address: edmarchiori@gmail.com (E. Marchiori).

<https://doi.org/10.1016/j.pulmoe.2020.04.011>

2531-0437/

© 2020 Sociedade Portuguesa de Pneumologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Severe rifampicin-induced thrombocytopenia in a patient with miliary tuberculosis



Tuberculosis is a potentially treatable infectious disease caused by *Mycobacterium tuberculosis* Complex. Pulmonary presentation is the most frequent, with non-pulmonary forms being more commonly observed in children and immunocompromised patients. Tuberculosis remains a major public health problem in Portugal, although its incidence has been decreasing (15,6 cases/100,000 inhabitants, according to the Portuguese National Statistical Institute, 2017). Newly diagnosed tuberculosis patients are empirically treated with a combination of four drugs:

isoniazid, rifampicin, pyrazinamide and ethambutol.¹ No antituberculous drug is free of risk, and the World Health Organization (WHO) recommends monitoring and reporting suspected or confirmed adverse drug reactions (ADR) caused by antituberculous drugs.² Most ADR are benign, result from inherent toxicity to the drug and can be minimized by dosage adjustment, exposure duration reduction or by vitamin supplementation.³ Less frequently, ADRs are immune-mediated and can present as cutaneous, hematological or systemic manifestations. These are rarer and more unpredictable, imposing both diagnostic and therapeutic challenges.⁴ Rifampicin, one of the most important antituberculous agents, is a well-tolerated and effective drug. Most frequent adverse reactions to rifampicin (Table 1) include gastrointestinal effects, cutaneous reac-