



LETTER TO THE EDITOR

Advantages and limitations of the ROX index



Dear editor

We have read with interest the study by Vega et al¹ published in the latest issue of the journal, where the authors propose the ROX index as a predictor of failure of high-flow nasal cannula (HFNC) therapy in patients with pneumonia due to SARS-CoV-2, and we would like to share some considerations on the advantages and disadvantages of using this index.

Non-invasive ventilatory support has gained relevance in recent years with the popularization of HFNC in patients with pneumonia. This therapy has been shown to be more effective than standard oxygen therapy and is recommended as first-line treatment for acute hypoxemic respiratory failure (AHRF).² These patients usually present dyspnea, hypoxemia, respiratory alkalosis, impaired gas exchange and consolidation images on chest tomography, similar to SARS-CoV-2 patients who also present fever and cough requiring a more advanced oxygen therapy.^{3,4}

By demonstrating its effects on gas exchange and respiratory mechanics, a possible delay in endotracheal intubation and invasive mechanical ventilation was quickly observed due to the possibility of masking the deterioration of the clinical picture. To avoid this situation, the ROX index (ratio of oxygen saturation as measured by pulse oximetry/ FiO_2 to respiratory rate) was proposed for patients with pneumonia and AHRF, and it showed accuracy in predicting HFNC failure at 12h of treatment (ROC 0.74 CI95% 0.64-0.84 $p < .002$), with <4.88 being the cut-off value associated with intubation (HR 0.273 CI95% 0.121-0.618 $p .002$).⁵

In the last 5 years, this index has been widely used due to its easy application at the bedside, which requires non-invasive variables for its measurement and can be evaluated at any time, even by non-medical health professionals. However, this same characteristic could cause small variations in its components to produce very dissimilar results. We must consider that the parameters to be evaluated can easily vary throughout the day or in different clinical situations (fever, mobilization, fatigue, pain, acidosis, hypotension). In addition, it could be considered as a static index that refers to a specific moment in time and not to the clinical evolution of the patient. Another disadvantage is that the index does not include the flow rate provided and it has been reported that

changes in the flow rate can modify the result of the therapy⁶ because it can generate a continuous pressure effect in the airway and favor the lavage of the dead space, increased end-expiratory volume and decreased respiratory rate and work of breathing. Due to these possible biases, other monitoring alternatives have been proposed, which we have discussed elsewhere.⁷ The role of lung ultrasonography (LUS) has also been mentioned as a tool to predict intubation: in addition to the evaluation of the excursion and diaphragmatic contraction, at bedside and non-invasively, LUS has proven the worsening of the disease in the presence of B lines pattern and the lack of aeration when dyspnea and hypoxemia were present.⁴

Vega et al demonstrated the usefulness of the ROX Index to guide the intubation decision in patients with COVID-19 pneumonia outside the ICU with a cut-off level <5.9 ,¹ however we suggest that the other parameters are not ignored, when taking decisions in scenarios of low vigilance, such as neurological deterioration, work of breathing, mental status alterations, agitation, drowsiness or stupor.

Funding

This manuscript did not present fundings of any kind.

Conflicts of interest

All authors declare no conflicts of interest.

Acknowledgments

None.

References

1. Vega ML, Dongilli R, Olaizola G, et al. COVID-19 pneumonia and ROX index: time to set a new threshold for patients admitted outside the ICU. *Pulmonology*. 2022;28(1):13–7. <https://doi.org/10.1016/j.pulmoe.2021.04.003>.
2. Lewis SR, Baker PE, Parker R, Smith AF. High-flow nasal cannulae for respiratory support in adult intensive care patients. *Cochrane Database Syst Rev*. 2021;3(3):CD010172. <https://doi.org/10.1002/14651858.CD010172.pub3>.

<https://doi.org/10.1016/j.pulmoe.2022.02.008>

2531-0437/© 2022 Sociedade Portuguesa de Pneumologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

3. Vega ML, Pisani L. Nasal high flow oxygen in acute respiratory failure. *Pulmonology*. 2021;27(3):240–7. <https://doi.org/10.1016/j.pulmoe.2021.01.005>.
4. Peixoto AO, Costa RM, Uzun R, Fraga AMA, Ribeiro JD, Marson FAL. Applicability of lung ultrasound in COVID-19 diagnosis and evaluation of the disease progression: a systematic review. *Pulmonology*. 2021;27(6):529–62. <https://doi.org/10.1016/j.pulmoe.2021.02.004>.
5. Roca O, Messika J, Caralt B, et al. Predicting success of high-flow nasal cannula in pneumonia patients with hypoxemic respiratory failure: the utility of the ROX index. *J Crit Care*. 2016;35:200–5. <https://doi.org/10.1016/j.jcrc.2016.05.022>.
6. Mauri T, Alban L, Turrini C, et al. Optimum support by high-flow nasal cannula in acute hypoxemic respiratory failure: effects of increasing flow rates. *Intensive Care Med*. 2017;43(10):1453–63. <https://doi.org/10.1007/s00134-017-4890-1>.
7. Gallardo A, Gigliotti C, Saavedra S, Zamarrón-López EI, Guerrero-Gutiérrez MA, Pérez-Nieto OR. Strategies for monitoring and predicting failure to high-flow nasal cannula therapy in the ED [published online ahead of print, 2021 Dec 24]. *Am J Emerg Med*. 2021;S0735-6757(21)01018–4.

A. Gallardo^{a,*}, E. Zamarrón-López^b,
E. Deloya-Tomas^c, O.R. Pérez-Nieto^c

^a *Sanatorio Clínica Modelo de Morón, Departamento de Kinesiología, Buenos Aires, Argentina*

^b *Hospital General Regional IMSS No. 6, Ciudad Madero, Unidad de Cuidados Intensivos, Tamaulipas, México*

^c *Hospital General San Juan del Río. Unidad de Cuidados Intensivos*

* Corresponding author.

E-mail address: adriankgallardo@gmail.com (A. Gallardo).

Received 18 February 2022; Accepted 23 February 2022

Available online 24 March 2022