



REVIEW

Issue 2 - “Update on adverse respiratory effects of indoor air pollution”. Part 2): Indoor air pollution and respiratory diseases: Perspectives from Italy and some other GARD countries



G. Sarno^a, I. Stanisci^a, S. Maio^a, S. Williams^b, E.M. Khoo^b, S.G. Diaz^c, E.V. Ponte^d, L.T.T. Lan^e, T. Soronbaev^f, D. Behera^g, S. Tagliaferro^a, S. Baldacci^a, G. Viegi^{a,*}

^a CNR Institute of Clinical Physiology (IFC), Via Trieste, 41, 56126 Pisa, Italy

^b International Primary Care Respiratory Group (IPCRG), 19 Armour Mews, Larbert FK5 4FF, Scotland, United Kingdom

^c Universidad Autónoma de Nuevo León, Faculty of Medicine and University Hospital “Dr. José Eleuterio González”, Regional Center of Allergy and Clinical Immunology, Av. Dr. José Eleuterio González 235, Mitras Centro, 64460 Monterrey, N.L., Mexico

^d Faculdade de Medicina de Jundiaí - Department of Internal Medicine, R. Francisco Teles, 250, Vila Arens II, Jundiaí SP, 13202-550, Brazil

^e University Medical Center, 217 Hong Bang, dist.5, Ho Chi Minh City 17000, Vietnam

^f Department of Respiratory Medicine, National Center for Cardiology and Internal Medicine, Togolok Moldo str., Bishkek 720040, Kyrgyzstan

^g Department of Microbiology, National Institute of Tuberculosis and Respiratory Diseases (NITRD), Sri Aurobindo Marg Near Qutub Minar, Mehrauli, New Delhi 110030, India

Received 20 March 2023; accepted 20 March 2023

Available online 19 May 2023

KEYWORDS

Household air pollution;
Chronic respiratory diseases;
Allergic diseases;
Epidemiology;
GARD;
Italy

Abstract

Objective: to synthesize the Italian epidemiological contribution to knowledge on indoor pollution respiratory impact, and to analyze the perspective of some GARD countries on the health effects of indoor air pollution.

Results: Italian epidemiological analytical studies confirmed a strong relationship between indoor air pollution and health in general population. Environmental tobacco smoke, biomass (wood/coal) fuel for cooking/heating and indoor allergens (house dust mites, cat and dog dander, mold/damp) are the most relevant indoor pollution sources and are related to respiratory and allergic symptoms/diseases in Italy and in other GARD countries such as Mexico, Brazil, Vietnam, India, Nepal and Kyrgyzstan.

Community-based global health collaborations are working to improve prevention, diagnosis and care of respiratory diseases around the world, specially in low- and middle-income countries, through research and education.

* Corresponding author at: CNR Institute of Clinical Physiology, Via Trieste, 41, 56126 Pisa, Italy.
E-mail address: viegig@ifc.cnr.it (G. Viegi).

Conclusions: in the last thirty years, the scientific evidence produced on respiratory health effects of indoor air pollution has been extensive, but the necessity to empower the synergies between scientific community and local administrations remains a challenge to address in order to implement effective interventions.

Based on abundant evidence of indoor pollution health effect, WHO, scientific societies, patient organizations and other members of the health community should work together to pursue the GARD vision of “a world where all people breathe freely” and encourage policy makers to increase their engagement in advocacy for clean air.

© 2023 Sociedade Portuguesa de Pneumologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Chronic respiratory diseases (CRDs) are among the most common non-communicable diseases worldwide. CRDs include chronic obstructive pulmonary disease (COPD), asthma, occupational lung diseases and pulmonary hypertension. The World Health Organization (WHO) Global Alliance against CRDs (GARD) vision is “a world in which all people breathe freely” and the aim of the WHO CRDs Programme is to support Member States, specially in low- and middle-income countries (LMICs), for reducing the toll of morbidity, disability and premature mortality of CRDs, specifically asthma and COPD.¹

Household air pollution (HAP) ranks 9th out of 69 risk factors linked to attributable deaths included in the Global Burden of Disease (GBD) project. HAP, caused by incomplete combustion of solid fuels and kerosene used for cooking/heating, was responsible for 3.2 million premature deaths in 2020 (over 237,000 in under 5 year-old children): 21% due to lower respiratory infections (LRI), 19% to COPD and 6% to lung cancer. HAP accounted for the loss of an estimated 86 million healthy life years in 2019, with the largest burden for women living in LMICs. Almost half of deaths due to LRI among under 5 year-old children are caused by PM (soot) from HAP.² Most deaths occur in three GBD Super Regions: South Asia (36%), sub-Saharan Africa (30%), and Southeast Asia, East Asia, and Oceania (28%).³

This paper is the fourth of the Series on “Air pollution and health”^{4–6} and is strictly linked to the companion paper on indoor pollution.⁷

Methods

There are two main sections, *an Italian perspective* and *the perspectives from other GARD countries*: the first summarizes results of the role of various indoor air pollutants on CRD-related outcomes in different Italian epidemiological studies; the second includes main results obtained in other GARD countries.

In Italy, epidemiological surveys on general population samples and school-children and adolescents were designed and conducted by the National Research Council (CNR), individually or in collaboration with other research groups, since the late 1970s. Health effects (e.g. allergic and respiratory symptoms/diseases, lung function tests, serum immunoglobulin E (IgE), exhaled nitric oxide (FeNO)) of indoor air

pollution in general population (adult and children) were assessed.^{8–21}

In particular, the Italian main results derive from the Po Delta (PD) and Pisa (PI) analytical prospective studies,^{12,22–28} “Italian Studies on Respiratory Disorders in Children and the Environment” (SIDRIA) study,^{15,29,30} “Epidemiologic study on environment and health in women” (SEASD) study,¹⁴ “Health Effects of School Environment” (HESE) study,^{31,32} “Geriatric study on health effects of air quality in nursing homes in Europe” (GERIE) study,¹⁸ the Palermo (PA) studies on children-adolescents,^{16,17} “Indoor and Outdoor Air Quality and Respiratory Health in Malta and Sicily” (RESPIRA) project^{19,20} and “Childhood Asthma and Environment Research” (CHASER) study.²¹

The perspectives from other GARD countries concern recent results from Mexico, Brazil, Vietnam, India, Nepal and Kyrgyzstan. An update on respiratory health impact of indoor air quality was performed through a PubMed Search for the period 2000–2022, focusing questionnaire and lung function tests. In addition, a part is dedicated to the community-based global health collaborations “Free Respiratory Evaluation and Smoke-exposure reduction by primary Health care Integrated groups” (FRESH AIR), “UK National Institute for Health Research, Global Health Research Unit on Respiratory Health” (RESPIRE) and “Global Health Respiratory Network” (GHRN).

Results

An Italian perspective

Results of CNR epidemiological surveys are summarized taking into account two different approaches of indoor air pollution assessment: qualitative evaluation from standardized questionnaires; quantitative evaluation (through active or passive samplers).

Indoor qualitative evaluation

In Italy (Table 1), it became clear in the early nineties that the fuel for cooking or for heating and that kind of heating are associated with increased prevalence rates of respiratory symptoms or impaired lung function. In PD males, significantly increased odds ratios (ORs) were found for: cough and dyspnea with bottled gas for cooking; cough and phlegm with stove for heating; cough with natural gas stove and fan or stove powered by other fuels.²³ In females, significantly increased ORs were found only for dyspnea with bottled gas

Table 1 Main studies carried out by CNR about the effects of indoor air pollution on respiratory health in Italy.

Author and year of publication	Population	Indoor setting	Exposure	Health outcome	Main results
Cooking and heating					
Viegi et al., 1991 (ref. 22)	381 subjects (8–19 year-old) living in Po River Delta area	Home	Cooking fuel and kind of heating	Chronic cough, attacks of shortness of breath with wheeze, wheeze, dyspnea, rhinitis and asthma	Among ETS exposed subjects, use of bottled gas for cooking than natural gas significantly associated with higher prevalence rates of wheeze (8%), dyspnea (11%) and diagnosis of asthma (11%).
Viegi et al., 1991 (ref. 23)	3289 subjects (8–64 year-old) living in Po River Delta area	Home	Cooking fuel and kind of heating	Chronic cough, chronic phlegm, dyspnea and lung function	<p>In males, significant associations between:</p> <ul style="list-style-type: none"> a) bottled gas for cooking (vs natural gas) and cough (OR 1.66; 95% CI 1.12–2.46) b) bottled gas for cooking (vs natural gas) and dyspnea (OR 1.81; 95% CI 1.15–2.85); c) stove for heating (vs central heating) and cough (OR 1.44; 95% CI 1.04–1.98); d) stove for heating (vs central heating) and phlegm (OR 1.39; 95% CI 1.01–1.91); e) stove fueled by natural gas (vs central heating fueled by natural gas) and cough (OR 1.54; 95% CI 1.10–2.35); f) fan or stove fueled other than by natural gas (vs central heating fueled by natural gas) and cough (OR 1.66; 95% CI 1.04–2.65). <p>In females, significant associations between:</p> <ul style="list-style-type: none"> a) bottled gas for cooking (vs natural gas) and dyspnea (OR 1.45; 95% CI 1.00–2.10) b) stove for heating (vs central heating) and dyspnea (OR 1.46; 95% CI 1.09–1.96); c) stove fueled by natural gas (vs central heating fueled by natural gas) and dyspnea (OR 1.58; 95% CI 1.07–2.34); d) stove or fan fueled other than by natural gas (vs central heating fueled by natural gas) and dyspnea (OR 1.73; 95% CI 1.12–2.67). <p>In females, use of stove than central heating significantly associated with lower values of FEF_{25–75}, MEF₅₀ and PEF.</p>

Table 1 (Continued)

Author and year of publication	Population	Indoor setting	Exposure	Health outcome	Main results
Viegi et al., 1992 (ref. 24)	3866 subjects (5–90 year-old) living in Pisa	Home	Cooking fuel and kind of heating	Cough, phlegm, wheeze, attacks of shortness of breath with wheeze	In non-smoking males, significant association between: a) bottled gas for cooking (vs natural gas) and cough (OR 2.82, 95% CI 1.12–7.10) In smoking males, significant or borderline associations between: a) bottled gas for cooking (vs natural gas) and cough (OR 1.39, 95% CI 0.98–1.96); b) bottled gas for cooking (vs natural gas) and phlegm (OR 1.42; 95% CI 1.00–2.01). In non-smoking females, significant associations between: a) stove or forced-air circulation (vs central heating) and wheeze (OR 1.54; 95% CI 1.07–2.21); b) stove or forced-air circulation (vs central heating) and shortness of breath with wheeze (OR 1.72; 95% CI 1.11–2.65).
SHS Paoletti et al., 1995 (ref. 25)	255 non-smoking subjects (8–20 year-old) living in the Po River Delta area	Home	SHS	Cough, phlegm, wheeze, attacks of shortness of breath with wheeze, asthma	In SHS exposed subjects than unexposed, significantly higher prevalence rates of cough (7.5%) and phlegm (5.9%).
Simoni et al., 2001 (ref. 26)	867 non-smoking women (8–73 year-old)	Home, school/work or elsewhere	SHS	Asthma (symptoms/diagnosis)	In non-smoking woman, SHS exposure vs no SHS exposure significantly associated with asthma (OR 1.90; 95% CI 1.1–3.4)
Simoni et al., 2003 (ref. 28)	120 elderly subjects (≥ 65 year-old) living in Pisa and Po River Delta area	Home	SHS	Acute respiratory symptoms	In SHS exposed subjects than unexposed, higher occurrence of acute respiratory symptoms both in winter (31 vs 29%) and summer (33 vs 16%).

Table 1 (Continued)

Author and year of publication	Population	Indoor setting	Exposure	Health outcome	Main results
Simoni et al., 2007 (ref. 14)	2195 never smoking women (mean age 47 years) living in Pisa, Po River Delta, Rome and Viterbo	Home/work	SHS to husband/partner, at workplace, in other places	Current dyspnea, any shortness of breath at rest, recent wheeze, recent attacks of shortness of breath with wheeze, asthma diagnosis/symptoms, obstructive lung diseases (diagnosis of asthma/chronic bronchitis/ emphysema), current cough/phlegm and rhino-conjunctivitis	In women, combined SHS exposure of husband/partner and at work vs no SHS exposure significantly associated with: a) current dyspnea (OR 1.61; 95% CI 1.20–2.16); b) any shortness of breath at rest (OR 2.81; 95% CI 1.83–4.30); c) recent wheeze (OR 1.71; 95% CI 1.04–2.82); d) recent attacks of shortness of breath with wheeze (OR 1.85; 95% CI 1.05–3.26); e) asthma diagnosis/symptoms (OR 1.50; 95% CI 1.09–2.08); f) obstructive lung diseases (OR 2.24; 95% CI 1.40–3.58); g) current cough/phlegm (OR 1.52; 95% CI 1.07–2.15); h) rhino-conjunctivitis (OR 1.48; 95% CI 1.13–1.94).
Cibella et al., 2011 (ref. 16)	2150 children (10–17 year-old) living in Palermo	Home	SHS	Current asthma and rhino-conjunctivitis	PAR% for current asthma attributable to SHS = 18.1%. PAR% for rhino-conjunctivitis attributable to SHS = 18.8%.
Ciluffo et al., 2022 (ref. 21)	179 asthmatic children (5–16 year-old) living in Palermo	Home	SHS	Uncontrolled asthma	In children, SHS exposure during pregnancy vs no SHS exposure during pregnancy associated with higher risk of uncontrolled asthma later in life (OR 3.82; 95% CI 1.11–13.06).

Table 1 (Continued)

Author and year of publication	Population	Indoor setting	Exposure	Health outcome	Main results
Particulate matter, nitrogen dioxide, formaldehyde and carbon dioxide					
Simoni et al., 2002 (ref. 27)	383 subjects (15–72 year-old) living in the Po River Delta	Home	NO ₂ -index* (high: > median value) and PM _{2.5} -index* (high: > median value)	Acute respiratory illnesses with fever, chronic bronchitis and/or asthmatic symptoms without fever, irritation without fever	<p>In winter, on the whole sample significant associations between:</p> <p>a) High NO₂-index (vs low) and acute respiratory illnesses with fever (OR 2.18; 95% CI 1.07–4.46);</p> <p>b) High NO₂-index (vs low) and chronic bronchitis and/or asthmatic symptoms without fever (OR 1.77; 95% CI 1.24–2.42).</p> <p>In winter, in non-smokers significant associations between:</p> <p>a) High NO₂-index (vs low) and acute respiratory illnesses with fever (OR 2.47; 95% CI 1.14–5.34);</p> <p>b) High NO₂-index (vs low) and chronic bronchitis and/or asthmatic symptoms without fever (OR 1.51; 95% CI 1.04–2.18);</p> <p>c) High PM_{2.5}-index (vs low) and chronic bronchitis and/or asthmatic symptoms without fever (OR 1.83; 95% CI 1.26–2.65).</p> <p>In summer, in non-smokers significant association between:</p> <p>a) High PM_{2.5}-index (vs low) and irritation without fever (OR 1.68; 95% CI 1.22–2.31)</p>
Simoni et al., 2003 (ref. 28)	120 elderly subjects (≥65 year-old) living in Pisa and Po River Delta area	Home	PM _{2.5} -index* (high: > median value)	Acute respiratory symptoms	<p>High PM_{2.5}-index exposure (vs low) associated with acute respiratory symptoms:</p> <p>a) 33 vs 27% in winter;</p> <p>b) 27 vs 21% in summer.</p>

Table 1 (Continued)					
Author and year of publication	Population	Indoor setting	Exposure	Health outcome	Main results
Simoni et al., 2004 (ref. 12)	1090 subjects (≥ 15 year-old) living in Pisa and Po River Delta area	Home	NO ₂ -index* (high: > median value) and PM _{2.5} -index* (high: > median value)	Acute respiratory illness, bronchitic and/or asthmatic symptoms without fever and without acute respiratory illness, increased PEF maximum amplitude, increased PEF diurnal variation	On the whole sample significant associations between: <ol style="list-style-type: none"> High NO₂-index (vs low) and acute respiratory illness with fever (OR 1.66; 95% CI 1.08–2.57); High PM_{2.5}-index (vs low) and acute respiratory illness with fever (OR 1.62; 95% CI 1.04–2.51); High PM_{2.5}-index (vs low) and bronchitic/asthmatic symptoms (OR 1.39; 95% CI 1.17–1.66); High PM_{2.5}-index (vs low) and increased PEF maximum amplitude (OR 1.38; 95% CI 1.24–1.54); High PM_{2.5}-index (vs low) and increased PEF diurnal variation (OR 1.37; 95% CI 1.23–1.53).
Simoni et al., 2010 (ref. 31)	654 children (10 year-old) living in five European countries (Siena and Udine, Italy; Reims, France; Oslo, Norway; Uppsala, Sweden; Århus, Denmark).	School	PM ₁₀ and CO ₂	Wheezing, dry cough at night, rhinitis and nasal patency	Schoolchildren exposed to elevated CO ₂ levels (> 1000 ppm) (vs low) showed a higher risk for: <ol style="list-style-type: none"> dry cough (OR 2.99; 95% CI 1.65–5.44); rhinitis (OR 2.07; 95% CI 1.14–3.73). Schoolchildren exposed to elevated PM ₁₀ levels (> 50 $\mu\text{g}\cdot\text{m}^{-3}$) (vs low) showed a lower nasal patency.
Cibella et al., 2015 (ref. 17)	303 children (10–17 year-old) living in Palermo	Home	NO ₂ (high: > 28 $\mu\text{g}/\text{m}^3$ in spring; > 41.4 $\mu\text{g}/\text{m}^3$ in winter)	Current asthma, wheeze, chronic phlegm and rhinoconjunctivitis	Children with current asthma were significantly more exposed to high level of indoor NO ₂ than low level (89% vs 11%); the same for children with wheeze episodes in the last 12 months (78% vs 22%), chronic phlegm (82% vs 18%), and rhinoconjunctivitis (66% vs 44%).

Table 1 (Continued)

Author and year of publication	Population	Indoor setting	Exposure	Health outcome	Main results
Bentayeb et al., 2015 (ref. 18)	600 elderly subjects (>65 year-old) living in seven European countries (Uppsala, Sweden; Warsaw, Poland; Athens, Greece; Reims, France; Arezzo, Italy; Århus, Denmark; Brussels, Belgium)	Nursing home	PM ₁₀ , PM _{0.1} , formaldehyde, nitrogen, NO ₂ and CO ₂ (high: > median value of the distribution of each pollutant)	FEV ₁ /FVC, breathlessness, cough, wheeze, COPD and exhaled carbon monoxide	Significant associations between: a) high PM _{0.1} (vs low) and FEV ₁ /FVC <70% (OR 8.16; 95% CI 2.24–29.3); b) high NO ₂ (vs low) and FEV ₁ /FVC <70% (OR 3.74; 95% CI 1.06–13.1); c) high PM ₁₀ (vs low) and breathlessness (OR 1.53; 95% CI 1.15–2.07); d) high PM ₁₀ (vs low) and cough (OR 1.73, 95% CI 1.17–10.3); e) high NO ₂ (vs low) and breathlessness (OR 1.58; 95% CI 1.15–2.20); f) high NO ₂ (vs low) and cough (OR 1.56; 95% CI 1.03–2.41); g) high PM _{0.1} (vs low) and wheeze (OR 2.82, 95% CI 1.15–7.02); h) high formaldehyde (vs low) and COPD (OR 3.49; 95% CI 1.17–10.3); i) high formaldehyde (vs low) and exhaled carbon monoxide (OR 1.25; 95% CI 1.02–1.55); j) high CO ₂ (vs low) and breathlessness (OR 1.68; 95% CI 1.32–2.15); k) high CO ₂ (vs low) and cough (OR 2.01; 95% CI 1.55–2.63).
Drago et al., 2018 (ref. 20)	73 children (10–15 year-old) living in Gela, Niscemi, Mazzarino and Butera (Sicily)	Home	PM _{2.5} mass concentration; fractions of Cd, Ce, La, and Tl in PM _{2.5} ⁺	Asthma symptoms in the last 12 months (wheezing/whistling in the chest and chest sounded wheezy during/after exercise or dry cough at night)	Asthma symptoms in the last 12 months significantly associated with: a) PM _{2.5} (OR 3.58; 95% CI 1.53–8.39); b) Cd (OR 1.53; 95% CI 1.01–2.31); c) Ce (OR 1.44; 95% CI 1.09–1.90); d) La (OR 1.49; 95% CI 1.11–2.01); e) Tl (OR 1.45 95% CI 1.08–1.95).

Table 1 (Continued)

Author and year of publication	Population	Indoor setting	Exposure	Health outcome	Main results
Allergens Simoni et al., 2005 (ref. 29)	20,016 children (6–7 year-old) and 13,266 adolescents (13–14 year-old) living in five areas of northern Italy (Turin, Milan, Mantua, Trent, and Emilia-Romagna), five areas of central Italy (Rome, Florence, Empoli, Siena, and Colli-ferro), and two areas of the south (Bari, Palermo).	Home	Moulds, dog and cat danders	Wheeze, asthma and cough/chronic phlegm	<p>In children, compared with no allergen exposure, significant associations between:</p> <ul style="list-style-type: none"> a) mould exposure in the first year of life and wheezing (OR 1.65; 95% IC 1.31–2.07); b) mould exposure in the first year of life and asthma (OR 1.80; 95% IC 1.41–2.30); c) mould exposure in the first year of life and cough/phlegm (OR 1.89; 95% IC 1.31–2.71); d) current mould exposure and wheeze (OR 1.62; 95% IC 1.22–2.15); e) dog exposure in the first year of life and wheezing (OR 1.44; 95% IC 1.01–2.05); f) dog exposure in the first year of life and cough/phlegm (OR 1.87; 95% IC 1.12–3.11). <p>In adolescents, compared with no allergen exposure, significant associations between:</p> <ul style="list-style-type: none"> a) mould exposure in the first year of life and wheezing (OR 1.56; 95% IC 1.15–2.11); b) mould exposure in the first year of life and asthma (OR 1.89; 95% IC 1.38–2.59); c) lifetime dog exposure and cough/phlegm (OR 3.69; 95% IC 1.66–8.21); d) dog exposure in the first year of life and cough/phlegm (OR 2.03; 95% IC 1.12–3.67).

Table 1 (Continued)

Author and year of publication	Population	Indoor setting	Exposure	Health outcome	Main results
Simoni et al., 2005 (ref. 30)	20,016 children (5–11 year-old) and 13,266 adolescents (12–17 year-old) living in five areas of northern Italy (Turin, Milan, Mantua, Trent, and Emilia-Romagna), five areas of central Italy (Rome, Florence, Empoli, Siena, and Colleferro), and two areas of the south (Bari, Palermo).	Home	Moulds/dampness	Asthma, rhino-conjunctivitis, wheeze and persistent cough/phlegm	<p>In children, compared with no allergen exposure, significant associations between:</p> <ul style="list-style-type: none"> a) mould/dampness exposure in the first year of life and asthma (OR 1.80; 95% CI 1.41–2.30); b) mould/dampness exposure in the first year of life and rhino-conjunctivitis (OR 1.46; 95% CI 1.17–1.82); c) mould/dampness exposure both in the first year of life/recently and wheeze (OR 1.98; 95% CI 1.47–2.66); d) mould/dampness exposure in the first year of life and cough/phlegm (OR 1.89; 95% CI 1.31–2.71); e) recently mould/dampness exposure and cough/phlegm (OR 1.86; 95% CI 1.19–2.91). <p>In adolescents, compared with no allergen exposure, significant associations between:</p> <ul style="list-style-type: none"> a) mould/dampness exposure in the first year of life and asthma (OR 1.89; 95% CI 1.38–2.59); b) mould/dampness exposure in the first year of life and wheeze (OR 1.56; 95% CI 1.15–2.11); c) mould/dampness exposure both in the first year of life/recently and rhino-conjunctivitis (OR 1.78; 95% CI 1.30–2.45).

Table 1 (Continued)

Author and year of publication	Population	Indoor setting	Exposure	Health outcome	Main results
Lombardi et al., 2010 (ref. 15)	20,016 children (6–7 year-old) living in five areas of Northern Italy (Turin, Milan, Mantua, Trent, Emilia-Romagna), four areas of Central Italy (Florence, Empoli, Siena, Colleferro) and two areas of Southern Italy (Bari, Palermo)	Home	Cat and dog danders	Current wheezing, current asthma and current rhinoconjunctivitis	Compared with no allergen exposure, significant associations between: <ol style="list-style-type: none"> cat exposure in the first year of life and current wheezing (OR 1.88; 95% CI 1.33–2.68); cat exposure in the first year of life and current asthma (OR 1.74; 95% CI 1.10–2.78); cat exposure in the first year of life and current rhinoconjunctivitis (OR 1.43; 95% CI 0.97–2.11).
Simoni et al., 2011 (ref. 32)	654 children (10 year-old) living in five European countries (Siena and Udine, Italy; Reims, France; Oslo, Norway; Uppsala, Sweden; Århus, Denmark).	School	Moulds (colony-forming units, cfu/m ³) and fungal DNA (cell equivalents, CE/g dust)	Past year dry cough at night, persistent cough, rhinitis, FEV ₁ and FVC	Children exposed to elevated viable mould levels (≥ 300 cfu/m ³) (vs low) showed higher risk of: <ol style="list-style-type: none"> past year dry cough at night (OR 3.10; 95% CI 1.61–5.98); persistent cough (OR 3.79; 95% CI 2.40–5.60); rhinitis (OR 2.86; 95% CI 1.65–4.95). 10 ⁴ CE/g dust increase of <i>Aspergillus/Penicillium</i> DNA was associated with wheeze (OR 2.14; 95% CI 1.05–4.36). 100 CE/g dust increase of <i>Aspergillus versicolor</i> DNA was associated with: <ol style="list-style-type: none"> wheeze (OR 1.48; 95% CI 0.99–2.21); cough (OR 1.75; 95% CI 1.17–2.62); rhinitis (OR 1.75; 95% CI 1.18–2.60). There were inverse associations of <i>Aspergillus versicolor</i> DNA with FVC (B –0.12; 95% CI –0.24–0.02) and <i>Streptomyces</i> DNA with both FEV ₁ (B –0.14; 95% CI –0.22–(–0.05)) and FVC (B –0.13; 95% CI –0.23–(–0.04))

Table 1 (Continued)					
Author and year of publication	Population	Indoor setting	Exposure	Health outcome	Main results
Cibella et al., 2011 (ref. 16)	2150 children (10–17 year-old) living in Palermo	Home	Moulds / dampness, <i>Dermatophagoides</i> mix, dog dander and cat dander	Current asthma and rhinoconjunctivitis	PAR% for current asthma attributable to: a) <i>Dermatophagoides</i> mix 19.8%; b) mould/dampness 7.6%; c) dog dander 5.3%; d) cat dander 7.2%. PAR% for rhinoconjunctivitis attributable to: a) <i>Dermatophagoides</i> mix 8.9%; b) mould/dampness 7.0%; c) dog dander 1.4%; d) cat dander 1.0%.
Ruggeri et al., 2017 (ref. 19)	125 children (10–14 year-old) living in a Southern Mediterranean (Sicily and Malta)	Home	Der p 1	Wheeze in the last 12 months, Der p 1-specific IgE level and FeNO	Der p 1 was positively correlated with both high Der p 1-specific IgE and FeNO in children with last 12 months wheeze.

OR: odds ratio.
95% CI: 95% confidence interval.
SHS: secondhand smoke.
NO₂: nitrogen dioxide.
RSP: Respirable Suspended Particulate Matter.
PM₁₀: particulate matter < 10 μm.
PM_{2.5}: particulate matter ≤ 2.5 μm.
PM_{0.1}: particulate matter <0.1 μm.
CO₂: carbon dioxide.
FEV₁: forced expiratory volume in one second.
FEF_{25–75}: forced mid-expiratory flow.
MEF₅₀: maximal expiratory flow at 50%.
PEF: peak flow.
eNO: exhaled Nitric Oxide.
PAR%: Population attributable risk, expressed as percent.
Der p 1: the major allergen of *Dermatophagoides pteronyssinus*.
FeNO: exhaled nitric oxide.
Cd: Cadmium.
Ce: Cerium.
La: Lanthanum.
Tl: Thallium.
IgE: Immunoglobulin E.
* NO₂-IndEx and PM_{2.5}-IndEx: daily indices of exposure to NO₂ and to PM_{2.5} computed as the product of weekly mean concentration of pollutant and the daily time of exposure (i.e. the time spent at home).
+ ORs and 95% CI for all variables along with the exposure to environmental tobacco smoke were evaluated as binary variable.
B: partial regression coefficient.

for cooking, stove for heating, natural gas stove, stove or fan powered by other fuels.²³ In PI males, significantly increased ORs were shown for: cough and phlegm associated with bottled gas for cooking in males; wheeze and shortness of breath with wheeze associated with the use of a stove or forced-air circulation in females.²⁴ In 8–19 years PD non-smokers, significantly higher prevalence rates of wheeze, dyspnea, diagnosis of asthma were found in subjects of both sexes using bottled gas for cooking and exposed to passive smoking.²²

The importance of secondhand smoke (SHS) exposure as a risk factor for respiratory diseases was also pointed out. Higher occurrence of cough and phlegm in young non-smokers and of acute respiratory symptoms in the elderly exposed to SHS were shown in PD and PI studies.^{25,28} In 867 PD non-smoking women, SHS was a significant risk factor for asthma.²⁶ In 2195 non-smoking women, living in PI, PD, Rome and Viterbo, SHS exposure both to husband/partner and at work was a significant risk factor for dyspnea, cough/phlegm and rhino-conjunctivitis, any shortness of breath at rest, wheeze and attacks of shortness of breath with wheeze, asthma diagnosis/symptoms, obstructive lung diseases (diagnosis of asthma/chronic bronchitis/emphysema).¹⁴ In 2150 PA schoolchildren, SHS was responsible for 18.1% of current asthma and 18.8% of rhino-conjunctivitis.¹⁶ Passive smoking exposure during pregnancy was related to a higher risk of uncontrolled asthma in a recent cohort study on PA asthmatic children.²¹

In the SIDRIA-2 study, carried out in 2002 in 12 areas of northern, central and southern Italy, mold exposure in the first year of life was positively associated with wheezing, asthma and rhino-conjunctivitis in children and adolescents, and with cough/phlegm only among children. Mould exposure in the previous year was a significant risk factor for wheeze, among children, and wheeze was most strongly related to early and current mould/dampness exposure; the same occurred in adolescents for rhino-conjunctivitis.^{29,30} In 2150 PA children, mould/dampness exposure was responsible of about 7% of current asthma and rhino-conjunctivitis cases.¹⁶

In the SIDRIA-2 study, dog exposure in the first year of life increased the risk for wheezing in children and for cough/phlegm in both children and adolescents. Lifetime dog exposure was significantly related to cough/phlegm in adolescents. Finally, cat exposure in the first year of life was significantly associated with current wheezing, asthma and rhino-conjunctivitis in children.^{15,29} In PA children, cat and dog exposures were responsible of 7.2% and 5.3% of current asthma cases, and of 1.4% and 1.0% of rhino-conjunctivitis cases, respectively.¹⁶

Indoor quantitative evaluation

In Italy (Table 1), the direct measurements of particulate matter < 2.5 μm ($\text{PM}_{2.5}$) and nitrogen dioxide (NO_2) were carried out in 140 PD households during winter and summer. The concentration of monitored pollutants was larger in winter than in summer. Significantly higher indoor NO_2 levels were found in the houses with gas-furnace heating and/or with gas water heater located inside the home. $\text{PM}_{2.5}$ was significantly higher in smokers' homes, and significantly related to the number of cigarettes smoked.³³ Only in winter, there were significant associations of bronchitic/

asthmatic symptoms with NO_2 and $\text{PM}_{2.5}$ values above the median. In summer, this association emerged for $\text{PM}_{2.5}$ only in non-smokers.²⁷ Moreover, in PD and PI adults, NO_2 and $\text{PM}_{2.5}$ exposures were associated with acute respiratory symptoms and mild lung function impairment.¹² In an elderly sample, the occurrence of acute respiratory symptoms was consistently higher in relation to the high respirable suspended PM (RSP) exposure, compared to low exposure.²⁸

In the GERIE study, for the first time, the relationships between indoor air quality, comfort parameters and respiratory morbidity among European elderly people living in nursing homes were investigated. The following associations were found: particulate matter < 0.1 μm ($\text{PM}_{0.1}$) with forced expiratory volume in one second (FEV_1)/ forced vitality capacity (FVC) < 70% and wheeze; PM_{10} and CO_2 with breathlessness and cough; NO_2 with FEV_1 /FVC < 70%, breathlessness and cough; formaldehyde with COPD and exhaled carbon monoxide. Pollutant effects were more pronounced in the case of poor ventilation.¹⁸

In the HESE study, schoolchildren exposed to CO_2 levels > 1000 ppm showed a higher risk for dry cough and rhinitis; nasal patency was significantly lower in those exposed to PM_{10} > 50 $\mu\text{g}\cdot\text{m}^{-3}$.³¹

PA adolescents, exposed to the highest indoor NO_2 concentrations, had increased frequency of current asthma, wheezing in the last 12 months, chronic phlegm, and rhino-conjunctivitis.¹⁷

In the RESPIRA project, assessment of respiratory health, determination of PM elemental composition and of selective markers of smoking were performed in 73 houses of adolescents living in Sicily (Gela, Niscemi, Mazzarino and Butera). When corrected for confounding factors, $\text{PM}_{2.5}$, Cadmium (Cd), Cerium (Ce), Lanthanum (La) and Thallium (Tl) were associated with increased probability of having respiratory symptoms.²⁰ Further, *Dermatophagoides pteronyssinus* (Der p 1) was positively correlated to both high Der p 1-specific IgE and Fractional exhaled nitric oxide (FeNO) in children with last 12 months wheeze.¹⁹

In the HESE study, an elevated concentration of moulds (≥ 300 cfu/m³) in the classroom was associated with higher risk of dry cough at night, persistent cough and rhinitis in the last 12 months. *Aspergillus/Penicillium* DNA was associated with wheeze, and *Aspergillus versicolor* DNA with wheeze, rhinitis and cough; there were inverse associations of *Aspergillus versicolor* DNA with FVC and *Streptomyces* DNA with both FEV_1 and FVC.³²

The perspectives from other GARD countries

A Mexican perspective

In Mexico (Table 2), 9–12 year-old children spend 85% of their time indoors.³⁴ Airborne pollen, indoor and outdoor pollutants may influence sick building syndrome (SBS).³⁵ In a study carried out in Monterrey (Nuevo Leon) 91% of subjects reported at least 1 SBS symptom, and fatigue was the most frequent (74.6%). *Quercus* and *Fraxinus* pollen were related with headaches, while humidity and NO_x were associated with eye and nasal symptoms.³⁵

In 5210 elementary schoolchildren of the El Paso Independent School District, ant and spider pest problems within the past year, pet dogs, fireplace heat, central air conditioning, humidifier use, and cooking with gas stoves were positively

Table 2 Main studies about the effects of indoor air pollution on respiratory health in some GARD countries.

Author and year of publication	Country	Population	Exposure	Health outcome	Main results
Cooking and heating					
Melson et al., 2001 (ref. 76)	Nepal	121 schoolchildren (11–17 year-old)	Open fire or burning stove without a flue	Asthma	The risk of asthma was higher among children exposed to smoky fuels (defined as living in a house with an open fire or burning stove without a flue) than for those not exposed (OR 2.2; 95% CI 1.0–4.5).
Regolado et al., 2006 (ref. 37)	Mexico	841 non-smoking women (≥ 38 year-old)	Cooking fuel	Phlegm, cough, wheezing and FEV ₁ /FVC	Among non-smoking women, use of biomass for cooking than natural gas significantly associated with higher prevalence rates of: a) bringing up phlegm all day (27.3%); b) wheezed in the past (46.1%); c) cough or phlegm all day (36.8%). Compared with those cooking with gas, current use of a stove burning biomass fuel was associated with reduced FEV ₁ /FVC (79.9 vs 82.8%).
Johnson et al., 2011 (ref. 66)	India	900 non-smoking women (≥ 30 year-old)	Biomass fuel	COPD	COPD prevalence was higher in biomass fuel users (vs clean fuel users) (OR 1.24; 95% CI 0.36–6.64)
da Silva et al., 2012 (ref. 48)	Brazil	1402 individuals	Indoor-Biomass and Outside-Biomass	Cough, wheezing and dyspnea	In adults (≥ 20 year-old), significant associations between: a) Indoor-Biomass (vs liquefied petroleum gas) and cough (OR 2.93; 95% CI 1.68–5.10); b) Indoor-Biomass (vs liquefied petroleum gas) and dyspnea (OR 2.59; 95% CI 1.32–5.09); c) Indoor-Biomass (vs liquefied petroleum gas) and wheezing (OR 2.33; 95% CI 1.25–4.38); d) Outside-Biomass (vs liquefied petroleum gas) and cough (OR 1.78; 95% CI 1.27–2.50); e) Outside-Biomass (vs liquefied petroleum gas) and dyspnea (OR 1.80; 95% CI 1.14–2.86);

Table 2 (Continued)

Author and year of publication	Country	Population	Exposure	Health outcome	Main results
					<p>f) Outside-Biomass (vs liquefied petroleum gas) and wheezing (OR 1.78; 95% CI 1.18–2.69). In young people (<20 year-old), significant associations between:</p> <p>a) Indoor-Biomass (vs liquefied petroleum gas) and cough (OR 1.64; 95% CI 1.16–2.31);</p> <p>b) Indoor-Biomass (vs liquefied petroleum gas) and dyspnea (OR 2.29; 95% CI 1.21–4.32);</p> <p>c) Indoor-Biomass (vs liquefied petroleum gas) and wheezing (OR 2.46; 95% CI 1.34–4.47);</p> <p>d) Outside-Biomass (vs liquefied petroleum gas) and wheezing (OR 1.77; 95% CI 1.15–2.72).</p>
Agrawal et al., 2012 (ref. 67)	India	156,316 subjects (20–49 year-old)	Cooking fuel	Asthma	In adult women, biomass and solid fuels (vs cleaner fuels) were associated with asthma (OR 1.26; 95% CI 1.06–1.49).
Kurmi et al., 2013 (ref. 72)	Nepal	1648 individuals (\geq 16 year-old)	Biomass smoke	FEV ₁ /FVC	Airflow obstruction (defined as FEV ₁ /FVC below the lower limit of normal) was significantly higher in subject exposed to the domestic biomass burning compared with those using liquid petroleum gas fuel (OR 2.06; 95% CI 1.16–3.67).
Bhat et al., 2013 (ref. 69)	India	214 children (<5 year-old)	Cooking fuel	ALRTI	Cooking fuel other than liquid petroleum gas (vs liquid petroleum gas) was associated with ALRTI (OR 3.58; 95% CI 1.23–10.45).
Acharya et al., 2015 (ref. 71)	Nepal	4802 children (<5 year-old)	Solid fuel	ARI	The risk of developing ARI was higher among children from the households using solid fuel in comparison with the children from households using cleaner fuel (OR 1.79; 95% CI 1.02–3.14).

Table 2 (Continued)					
Author and year of publication	Country	Population	Exposure	Health outcome	Main results
Sinha et al., 2017 (ref. 73)	India	1200 adults (≥ 30 year-old)	Biomass fuel	COPD	Biomass fuel use resulted strongly associated with COPD (AOR 2.64; 95% CI 1.48–4.71)
Svendsen et al., 2018 (ref. 36)	Mexico	5210 children (9–11 year-old)	Fireplace heat, central air conditioning, humidifier use and cooking with gas stoves	Allergy and asthma	<p>Allergy prevalence:</p> <ul style="list-style-type: none"> a) 21.1% in presence of fireplace heat; b) 84.8% in presence of central air conditioning; c) 23.1% in presence of humidifier use; d) 81.6% in presence of cooking with gas stoves. <p>Ever asthma prevalence:</p> <ul style="list-style-type: none"> a) 22.6% in presence of fireplace heat; b) 85.7% in presence of central air conditioning; c) 23.1% in presence of humidifier use; d) 81.2% in presence of cooking with gas stoves. <p>Current asthma prevalence:</p> <ul style="list-style-type: none"> a) 22.3% in presence of fireplace heat; b) 86.3% in presence of central air conditioning; c) 25.1% in presence of humidifier use; d) 80.5% in presence of cooking with gas stoves.
Ramírez-Venegas et al., 2018 (ref. 38)	Mexico	1333 women (>35 year-old)	Biomass smoke	Cough, phlegm, wheezing, dyspnea, FEV ₁ , FVC, FEV ₁ /FVC and SaO ₂	<p>The prevalence of respiratory symptoms was significantly higher in women with biomass smoke exposure compared with those without exposure:</p> <ul style="list-style-type: none"> a) cough (24%); b) phlegm (29%); c) wheezing (25%); d) mMRC1 (27%); e) mMRC2 (16%).

Table 2 (Continued)					
Author and year of publication	Country	Population	Exposure	Health outcome	Main results
					<p>The spirometry and oxygen saturation values were significantly lower in women with greater biomass exposure levels:</p> <ul style="list-style-type: none"> a) at 100–199 h/years of exposure FEV₁ was 2.09 L ± 0.50 L; b) at 100–199 h/years of exposure FVC was 2.57L±0.52 L; c) at 100–199 h/years of exposure FEV₁/FVC was 80±8; d) at 100–199 h/years of exposure SaO₂ was 92.81%±2.9%; e) at 200–600 h/years of exposure FEV₁ was 1.86L±0.50 L; f) at 200–600 h/years of exposure FVC was 2.34L±0.55 L; g) at 200–600 h/years of exposure FEV₁/FVC was 79±6; h) at 200–600 h/years of exposure SaO₂ was 91.9%±2.2%.
Fernandes et al., 2018 (ref. 46)	Brazil	996 subjects (≥18 year-old)	Biomass smoke	Asthma	<p>In individuals with asthma, exposure to smoke from wood stoves vs no exposure significantly associated with:</p> <ul style="list-style-type: none"> a) lack of asthma control (OR 1.75; 95% CI 1.20–2.56); b) greater severity (OR 1.46; 95% CI 1.04–2.05).
Naz et al., 2018 (ref. 70)	Nepal	17,780 children (<5 year-old)	Cooking fuel	Mortality	<p>The use of cooking fuel was associated with total under-five mortality (OR 2.19; 95% CI 1.37–3.51), with stronger associations evident for sub-group analyses of neonatal mortality (OR 2.67; 95% CI 1.47–4.82)</p>

Table 2 (Continued)

Author and year of publication	Country	Population	Exposure	Health outcome	Main results
Baggio et al., 2019 (ref. 44)	Brazil	22,485 individuals (>18 year-old)	Solid fuels	Bronchitis/asthma	In women, solid fuels vs clean fuels associated with higher risk of bronchitis/asthma (OR:1.43).
Brakema et al., 2019 (ref. 81)		Kyrgyzstan	392 individuals (>18 year-old)	PM _{2.5}	COPD The highest PM _{2.5} exposure (vs the lowest exposure) associated with COPD (OR 3.17, 95% CI 1.01–9.49).
Rao et al., 2022 (ref. 68)	India	4662 individuals (>30 year-old)	Biomass fuel (wood, dung, crop) and mosquito coil	Poor respiratory health	Poor respiratory health was associated with: a) use of a combination of all three biomass fuel types (AOR 1.69; 95% CI 1.13–2.54); b) use of mosquito coil more than or equal to 5 days per week (AOR 1.43; 95% CI 1.04–1.99).
SHS Melsom et al., 2001 (ref. 76)	Nepal	121 schoolchildren (11–17 year-old)	SHS	Asthma	Asthma was associated with cigarette smoking by two or more family members (OR 1.9; 95% CI 1.0–3.9).
Gupta et al., 2006 (ref. 75)	India	62,109 adults	SHS	Asthma	Asthma was associated with: a) SHS exposure during childhood only (OR 1.38; 95% CI 1.08–1.75); b) SHS exposure both during childhood and adulthood (OR 1.22; 95% CI 1.04–1.43).
Dahal et al., 2009 (ref. 77)	Nepal	6056 children (<5 year-old)	SHS	ARI	The risk of ARI symptoms was higher among children born to mothers who smoked more frequently than their counterparts (AOR 1.41; 95% CI 1.02–1.96).
Schvartsman et al., 2013 (ref. 53)	Brazil	105 children (6–23 months-old)	SHS	Wheezing	On the whole sample significant associations between: a) from 1 to 7 cigarettes and wheezing (OR 6.98; 95% CI 1.73- 28.20); b) from 8 to 90 cigarettes and wheezing (OR 32.15; 95% CI 5.85–176.61)

Table 2 (Continued)

Author and year of publication	Country	Population	Exposure	Health outcome	Main results
Fernández-Plata et al., 2016 (ref. 39)	Mexico	3187 schoolchildren (8–17 year-old)	SHS	Respiratory symptoms (cough, wheezing, phlegm and dyspnea), respiratory infections (common cold, sore throat and bronchitis), FEV ₁ and FVC	OR of between 1.4 and 2.3 predicting respiratory symptoms in those exposed to passive smoking, exhibiting a trend to increase with the number of persons smoking inside the home; for recent respiratory infections, OR were between 1.2 and 1.5. Levels of FEV ₁ and FVC in individuals exposed to passive smoking were 6.8 and 14.1 ml, respectively, below those of non-exposed children.
Freire et al., 2016 (ref. 52)	Brazil	120 subjects (30–50 year-old)	SHS	FEV ₁	FEV ₁ % predicted was significantly lower in the passive smoking group when compared with the control group (93.41 ± 14.90 vs 101.9 ± 9.81).
Fernandes et al., 2018 (ref. 54)	Brazil	375 adolescents (13-14 year-old)	SHS	Wheezing	Wheezing was associated with more than 20 cigarettes per day smoked by parents or other household members (OR 3.6; 95% IC 1.2–10.7).
Thapa et al., 2016 (ref. 78)	Nepal	198 children (<5 year-old)	SHS	ARI	ARI was associated with: a) non-exclusively breastfed children with the presence of anyone smoking at their house (AOR 4.8, 95% CI 1.7–13.2), b) smoking in presence of children (AOR 6.4, 95% CI 1.1–38.3).
Schivinski et al., 2017 (ref. 51)	Brazil	78 schoolchildren (6–14 year-old)	SHS	PEF	In SHS exposed children there was lower PEF (4.194 L/s ± 1.278 vs 5.064 L/s ± 1.604) than in those unexposed.
Sinha et al., 2017 (ref. 73)	India	1200 adults (≥ 30 year-old)	SHS	COPD	The risk of COPD was higher among subjects exposed to SHS than for those not exposed (AOR 7.9; 95% CI 3.32–13.18).

Table 2 (Continued)

Author and year of publication	Country	Population	Exposure	Health outcome	Main results
Rivera et al., 2021 (ref. 40)	Mexico	536 children (6–8 year-old)	SHS and PM _{2.5}	Wheeze	Among children born to mothers exposed to SHS in pregnancy when compared with unexposed mothers, significant association between higher prenatal and early life PM _{2.5} exposure and higher cumulative risk ratios of: a) ever wheeze (RR 3.76; 95% CI 1.41–10.0 per 5 µg/m ³), b) current wheeze in the past year (RR 7.9; 95% CI 1.5–41.6 per 5 µg/m ³)
Allergens Lâm et al., 2014 (ref. 62)	Vietnam	684 subjects (23–72 year-old)	Any mite, cockroach, any animal and to any allergen	AR	AR were strongly associated with sensitisation to: a) any mite ($p = 0.009$); b) cockroach ($p = 0.001$); c) any animal ($p = 0.001$); d) any allergens ($p < 0.001$). AR symptoms last 12 months were strongly associated with sensitisation to: a) any mite ($p = 0.003$); b) cockroach ($p = 0.002$); c) any animal ($p < 0.001$); d) any allergens ($p < 0.001$). Persistent allergic rhinitis was significantly associated with sensitisation to: a) cockroach ($p < 0.001$); b) any animal ($p < 0.001$); c) any allergens ($p = 0.034$).
Álvarez-Chávez et al., 2016 (ref. 41)	Mexico	18 sensitized children (3–6 year-old)	Dust	Specific sensitivity (IgE) to <i>Dermatophagoides pteronyssinus</i>	Significant association found was between the presence of dust and sensitivity with the allergen <i>Dermatophagoides pteronyssinus</i> (dust mite) (Kappa value: 0.44; $p = 0.01$).

Table 2 (Continued)					
Author and year of publication	Country	Population	Exposure	Health outcome	Main results
Fernandes et al., 2018 (ref. 54)	Brazil	375 adolescents (13–14 year-old)	Pets	Wheezing	Contact with animals in the first year of life (dogs, cats, other furry animals, or birds) was associated with wheezing (OR 3.2; 95% IC 1.1–8.8).
Svensden et al., 2018 (ref. 36)	Mexico	5210 children (9–11 year-old)	Ant, spider and dog	Allergy and asthma	Allergy prevalence: a) 25.1% in presence of ant; b) 11.5% in presence of spider; c) 54.3% in presence of dog. Ever asthma prevalence: a) 25.0% in presence of ant; b) 12.1% in presence of spider; c) 53.9% in presence of dog. Current asthma prevalence: a) 25.2% in presence of ant; b) 12.7% in presence of spider; c) 53.6% in presence of dog.
Segura-Medina et al., 2019 (ref. 42)	Mexico	42 patients with asthma (18–76 year-old)	Moulds	ACT, FEV ₁ and FVC	Concentrations of some moulds were negatively associated with parameters of asthma control in male subjects, but not in female patients; significant Spearman correlation coefficient in male were: a) –0.72 for the relationship between <i>Aspergillus fumigatus</i> and FVC; b) –0.70 for the relationship between <i>Aspergillus fumigatus</i> and FEV ₁ ; c) –0.69 for the relationship between <i>Aureobasidium pullulans</i> and FEV ₁ ; d) –0.65 for the relationship between <i>Stachybotrys chartarum</i> and FEV ₁ /FVC; e) –0.65 for the relationship between <i>Alternaria alternata</i> and FEV ₁ ; f) –0.60 for the relationship between <i>Alternaria alternata</i> and FEV ₁ /FVC;

Table 2 (Continued)					
Author and year of publication	Country	Population	Exposure	Health outcome	Main results
					g) -0.80 for the relationship between <i>Cladosporium cladosporioides</i> 2 and ACT; h) -0.60 for the relationship between <i>Cladosporium herbarum</i> and FEV ₁ ; i) -0.74 for the relationship between <i>Epicoccum nigrum</i> and ACT.
Neto et al., 2019 (ref. 55)	Brazil	878 children (9–12 year-old)	Cat	Asthma	Having a cat during the first year of life was a risk factor associated with asthma (OR 1.73; 95% IC 1.07–2.78). Compared with no allergen exposure, significant associations between: a) dampness and asthma (OR 5.21; 95% CI 1.23–22.10); b) pets and asthma (OR 6.77; 95% CI 1.76–25.99).
Deka et al., 2022 (ref. 74)	India	150 patients (3–12 year-old)	Pets and dampness	Asthma	
<p>FEV₁: forced expiratory volume in the first second. FVC: forced vital capacity. ALRTI: acute lower respiratory tract infections. ARI: acute respiratory tract infections. COPD: chronic obstructive pulmonary disease. AOR: adjusted odds ratio. SaO₂: oxygen saturation. mMRC: Modified Medical Research Council Dyspnea Scale. OR: odds ratio. 95% CI: 95% confidence interval. SHS: secondhand smoke. PM_{2.5}: particulate matter $\leq 2.5 \mu\text{m}$. RR: cumulative risk ratios. ACT: Asthma Control Test. AR: allergic rhinitis. PEF: peak flow.</p> <p>Poor respiratory health: (i) individual with symptoms of chronic bronchitis, namely cough with expectoration for three or more months; for two or more years (ii) wheezing or chest tightness/ breathlessness in the morning and positive response to either of the three questions; previous diagnosis of asthma, use of medication for asthma or history of attack of asthma in the previous 12 months that made the respondent feel short of breath.</p>					

associated with both allergy and asthma prevalence (ever and current). Lung function decreased among children who lived in homes with reported cockroach pest problem in the past year without concurrent use of pesticides.³⁶

In 841 non-smoking ≥ 35 year-old women in the village of Solis (near Mexico City), compared with those cooking with gas, current use of a stove burning biomass fuel was associated with increased reporting of phlegm and reduced FEV₁/FVC.³⁷ In > 35 year-old females in a suburban area of Mexico City, spirometry and oxygen saturation values were significantly lower with higher biomass exposure levels. Cough, phlegm, wheezing and dyspnea were significantly more frequent in women exposed to biomass.³⁸

In 8–17 year-old schoolchildren from Mexico City and its metropolitan area ($n = 3187$), passive smoking was associated with a higher frequency of self-reported respiratory symptoms and respiratory infections. Levels of FEV₁ and FVC in children exposed to passive smoking were reduced: such values decreased with increasing number of smokers at home and higher outdoor ozone levels.³⁹ A study which involved 536 mother-child couples showed significant associations of higher prenatal and early life PM_{2.5} exposure with higher cumulative risk ratios of ever wheeze and current wheeze in the past year only among 6–8 year-old children born to mothers exposed to SHS in pregnancy.⁴⁰

A significant association between dust and sensitivity to dust mite was found in homes of sensitized children in Hermosillo (Sonora).⁴¹ In 42 adult asthma patients in Mexico City, mould burden was negatively associated with parameters of asthma control only in males.⁴²

Identification and quantification of persistent indoor organic pollutants were carried out in winter within three bus lines of Leon (Guanajuato) in order to determine their possible origin, the differences in the levels of contamination between routes, and the potential risk to the health of the users. Xylenes were the most representative organic pollutants with the potential to generate a higher risk to bus-users' health (hazard quotient: 6.4–9.8), without dismissing the potential danger of other pollutants.⁴³

A Brazilian perspective

In Brazil (Table 2), wood smoke is a major source of indoor air pollution, mainly among non-affluent people in rural areas.⁴⁴ In Bahia state, childhood exposure to indoor wood smoke was reported by 84% adult rural-born asthmatics vs only 27% urban-born subjects;⁴⁵ 1.7% of asthmatic urban dwellers reported current use of wood stoves for cooking.⁴⁶ In the north and northeast region, residential wood burning was significantly associated with higher risk of bronchitis/asthma only in women.⁴⁴ The PURE study, a multinational 10 year cohort study on subjects from 22 urban communities and 42 rural areas, showed that Brazilians using solid fuel for cooking had higher cardiorespiratory mortality rate, compared to users of electric or gas stoves.⁴⁷ In 1402 subjects living in Joao Camara, significantly increased ORs for cough, wheezing and dyspnea emerged in adults exposed to indoor stove and outside bonfire, compared to liquefied petroleum gas. Both non-smokers exposed to biomass and smokers exposed to gas had decreased %predicted-FEV₁ and FEV₁/FVC, as compared to non-smokers exposed to gas.⁴⁸

Policies increasing access to clean fuel strongly reduced the utilization of wood fire during the last five decades.⁴⁹

However, the interruption of the exposure does not avoid completely its late effects on respiratory health: in fact, previously exposure to indoor wood smoke predicts later uncontrolled asthma symptoms and lung function impairment in subjects no longer exposed, in comparison to never exposed ones.⁴⁶

Tobacco-free policy contributed to the decreasing prevalence of smoking during the last few decades, but second-hand smoking remains an important source of indoor air pollution.⁵⁰ It is associated with central and peripheral airway obstruction in children and adolescents.⁵¹ In adults, passive smoking is associated with impaired mucociliary clearance: exposure load correlates with worse hemodynamic parameters, reduced lung function and autonomic nervous system dysfunction.⁵² The probability of wheezing increased with the number of cigarettes smoked indoors in 105 children (6–23 months of age).⁵³

In 375 Belo Horizonte adolescents, asthma symptoms were related to contact with animals during the first year of life and over 20 cigarettes per day smoked by parents or other household members.⁵⁴ In 878 9–12 year-old children in Passo Fundo (Rio Grande do Sul), having an indoor cat during the first year of life was associated with asthma.⁵⁵

PM is another source of indoor air pollution in large urban centers. In 2018, a strong correlation between indoor and outdoor concentrations of PM₁₀ and PM_{2.5} in a nursing home in São Paulo City over thirty days of sequential sampling was observed.⁵⁶ During the sampling period, indoor PM concentration exceeded the 2005 WHO guidelines.⁵⁶ In the same year, high indoor concentrations of fine PM were observed in most studied residences in São Paulo metropolitan area, with indoor PM_{2.5} exceeding WHO recommendations in 43% of the residences; the highest values occurred in houses near heavy traffic areas during non-precipitating days.⁵⁷ However, recent studies relating indoor concentration of PM and respiratory health are lacking.

A Vietnamese perspective

In Vietnam (Table 2), $> 64\%$ people live in rural areas with less accessibility to cleaner fuels. Energy Information Administration reported that about 25% of the total Vietnam's energy consumption was from solid biomass fuels and wastes: nearly 60% of biomass was consumed by households, mainly in rural areas.⁵⁸

SHS is an important source of HAP: male smoker prevalence is very high (45.3%).

Despite the public places' bans, SHS prevalence among adults ≥ 15 years remained high (89.1% in Bars/Cafes/Tea shops and 80.7% in restaurants, 37.9% in government offices, and 30.9% in universities).⁵⁹

House ventilation is poor, especially in crowded city. People live very close to roadways and busy streets, which leads to increase PM and NO₂ inside the houses. Recent studies, measuring outdoor and indoor air quality in Hanoi, were focused on particle number and particle mass concentrations, which showed the influence of traffic and construction activities in both outdoor and indoor environment.^{60,61}

In South-East Asia, high temperature and humidity are an ideal habitat for mites and cockroaches.^{62,63} In Vietnam, the humidity and dampness are high, especially in Red River and Mekong delta.

Thus, moulds, bacteria, insects and mites grow rapidly and are linked to asthma, chronic rhinosinusitis, hypersensitivity pneumonitis and exacerbations of underlying respiratory diseases.⁶⁴

In urban and rural environments, there are many allergens from house dust mites, cockroaches, and mouse infestations, contributing to respiratory diseases such as wheeze, asthma attacks, asthma, and allergic rhinitis, as emerged in 864 adults living in northern Vietnam.⁶²

Many domestic behaviors and practices are associated with increased HAP: incense burning, poor ventilation and keeping moto-bikes in small houses, high concentrations of CO₂ and volatile organic compounds (VOCs).

To reduce HAP in rural areas, use of cleaner biomass burning stoves, chimney for kitchen and gas to replace biomass fuels were incentivized. However, traditional way of cooking is still mostly used in remote areas.

In urban regions, better ventilation and use of air cleaners were encouraged, but the problem of residential overcrowding is very difficult to solve. Reducing the prevalence of smoking is a major goal, but it is time consuming.

The Projet Interuniversitaire Cible (PIC), between universities of Vietnam and Belgium from the year 2014 to 2023 aiming at reducing HAP, is being carried out with some preliminary positive results.⁶⁵

A perspective from India and Nepal

In a study conducted among 900 \geq 30 year-old non-smoking women, from 45 rural villages of Tiruvallur district (Tamil Nadu, India) (Table 2), COPD prevalence was higher in biomass fuel users than in clean fuel users.⁶⁶ The same association emerged for asthma in women within a nationwide large-scale cross-sectional survey on 156,316 subjects aged 20–49 years.⁶⁷ In a survey on \geq 35 year-old adults in 5 villages of Tiruvallur district, combined use of all three biomass fuel types (wood, dung and crop) and use of mosquito coil \geq 5 days per week were associated with poor respiratory health.⁶⁸ In 214 $<$ 5 year-old Indian children, cooking fuel other than liquid petroleum gas was found to be significant risk factors for acute LRI.⁶⁹ From 2001 to 2011, use of cooking fuel was associated with total mortality in $<$ 5 years children from Nepal, with stronger associations evident for \leq 28 days infants mortality.⁷⁰ Associations were found also with acute respiratory tract infections (ARI).⁷¹

In a Nepalese population sample, prevalence of airflow obstruction was significantly higher in subject exposed to the domestic biomass burning, compared with those using liquid petroleum gas fuel.⁷² In 1200 \geq 30 year-old adults in Mehrauli, a large urban community in South Delhi (India), SHS and biomass fuel use were found among the risk factors associated with COPD.⁷³

In 150 clinically diagnosed 3–12 year-old asthmatics, damp environment, pets and passive smoking were significant risk factors for asthma.⁷⁴

In a multicentric population study carried out in India, SHS exposure during childhood is an important risk factor for asthma and respiratory symptoms in non-smoking adults: a higher risk of having asthma emerged for exposures in childhood only or both in childhood and adulthood.⁷⁵ In a Nepalese study, children asthma was associated with cigarette smoking by two or more family members and domestic use of smoky fuels.⁷⁶ In a Nepalese study carried out on 6056

children, maternal smoking had a significant effect on children's ARI symptoms, mainly among those born to mothers who smoked more frequently.⁷⁷ More recently, in a study carried out in Kathmandu, non-exclusive breastfeeding (with respect to exclusive one) increased the effect of household passive smoking on ARI.⁷⁸

A perspective from Kyrgyzstan

In Kyrgyzstan (Table 2), indoor pollution is a major risk factor among highlanders who use only biomass for cooking and heating, in poorly ventilated rooms. High level of indoor air pollution may cause respiratory symptoms among highlanders.^{79,80}

In a study conducted in a highland and a lowland rural setting, COPD prevalence and indoor PM_{2.5} exposure was highest in the highlands, and were independently associated.⁸¹

The FRESH AIR research program found biomass use for heating and cooking by 71.2% and 52.0% of a sample of COPD patients, respectively;⁸² therefore, the health impact of reducing biomass exposure could be considerable. Among 99 adults and children living in rural communities, after implementing improved cookstoves/heaters, significant reductions of PM_{2.5} concentration (65%) and of daily respiratory symptoms and chest infections were found.⁸³

A perspective from the International Primary Care Respiratory Group (IPCRG)

The IPCRG is a clinically-led charitable organization for improving prevention, diagnosis and care of respiratory diseases in community settings around the world through research and education. Since 2011, in collaboration with multiple partners, IPCRG has been engaging with stakeholders in eight resource-limited settings in three continents to measure and to co-create evidence-informed programmes addressing exposure to poor indoor air and its impact on respiratory health. Where evidence about relevant interventions existed, implementation science methods were applied to test its relevance, impact and generalizability in different settings. Where it was lacking, knowledge gaps were identified and tacit colloquial and local evidence elicited to enrich understanding and to inform the development of new interventions.⁸⁴

An IPCRG's FRESH AIR protocol was developed in Vietnam and in rural Uganda, contributing to an understanding that COPD may start early in life, and prevalence is highest in men and women aged 30–39, due to both high male smoking rates and also exposure to indoor biomass.^{85,86} RESPIRE, a Global Health Research Unit at the University of Edinburgh funded by the National Institute for Health Research (NIHR), aims to reduce the impact and number of deaths caused by respiratory diseases in Asia. A 2019 systematic review by RESPIRE found that the global prevalence of COPD among 30–79 year-old people was between 7.6%–10.3% depending on the definition used, and that smoking, biomass exposure and occupational exposure to dust or smoke were all substantial risk factors for COPD.⁸⁷ RESPIRE and FRESH AIR researchers subsequently collaborated to review prevalence survey methodology for CRDs in LMICs to inform new surveys in Asia, finding that methods to map and measure COPD are more reliable than asthma.⁸⁸

An adapted FRESH AIR protocol was applied in four countries with limited resources (Vietnam, Crete, Uganda and the Kyrgyz Republic), as part of a European Commission Horizon 2020 programme 2015–2019.⁸⁹ It included:

- Structured rapid assessment highlighting three key factors that needed to be addressed: 1. perceived CRD identity when systems and education have been primarily about respiratory infection leading to inaccurate attribution of symptoms; 2. beliefs about causes of CRD: 65% strongly agreed that tobacco smoking caused symptoms, but only 19% for household air pollution; and other perceived causes included witchcraft and a hot-cold disbalance and 3. norms and social structures for example “real men smoke”.⁹⁰
- Capacity building for health care providers (HCPs) to deliver Very Brief Advice (VBA) for quitting tobacco, pulmonary rehabilitation and community education about the harms of indoor air pollution from burning biomass fuels in the home.⁹¹
- Use of culturally appropriate engagement strategies including the use of photography and film.

It was concluded that contextualizing the programmes is feasible, acceptable and effective, and increases their sustainability.⁹¹

An estimated 2.6–3 billion people rely on solid fuels for cooking or heating increasing their risk of harm to their respiratory health. The impact of financial austerity may increase this use.⁹² Therefore, accelerating access to cleaner solutions was a target of many studies from the perspective of health, environment, climate and gender equity: yet, the results tended to be disappointing. IPCRG coordinates the Global Health Respiratory Network (GHRN), established in 2018, focusing on the poorest communities embracing use of clean cookstoves and clean fuel.⁹³ Two studies explored the barriers to uptake and found that funds, knowledge and beliefs about the innovation of solid fuel cookstoves and clean fuels were common barriers. External policy and incentives are also important for clean fuels.⁹⁴ Meanwhile, “disconnects” for example between solid fuel usage and perceived health impact and gender-related disconnect need to be understood.⁹⁵ Co-developing local solutions with the community was important for success, including using sensory perceptions of mostly invisible air pollution to improve understanding. The Cleaner Cookstove Implementation Tool and the Clean Fuel Implementation Tool were proposed.⁹⁴

FRESH AIR also revealed the socioeconomic impact of CRD in low-income settings: in particular, the “presenteeism” phenomenon where people attend school and work but their productivity drops; the desire among communities to improve the quality of the air they breathe. Where system-level changes are needed, making the case for change in resource-limited settings can be a major barrier.

Discussion

An Italian perspective

The Italian epidemiological analytical studies confirmed a strong relationship between indoor air pollution and health

status in the general population. Cooking and heating fuel, as well as SHS, were related to respiratory and allergic symptoms/diseases; exposures to NO₂ and PM_{2.5} were associated with acute and chronic respiratory effects (symptoms, impaired lung function); allergen exposure was related to respiratory/allergic outcomes depending on allergen type and exposure period, with the strongest effects for early-life exposure.

In Italy, the GARD (GARD-I) initiative was launched in 2009 at the Ministry of Health which assumed the role of national coordinator. The overall objective of GARD-I is to reduce incidence, morbidity and mortality of respiratory diseases through an integrated and intersectoral approach aimed at their prevention, including the fight against the main modifiable risk factors.⁹⁶ The working group “Project n.1 Prevention program for schools of indoor risks for respiratory and allergic diseases” produced a document on air quality in schools and risks for respiratory and allergic diseases in Italy, summarizing also the main steps to be taken to reduce indoor risks in school environments and warranting students’ well-being.⁹⁷

Italian research groups participated in the SINPHONIE (Schools Indoor Pollution and Health – Observatory Network in Europe) project, performed by a consortium of 38 partners from 25 countries, contributing to release guidelines for healthy environments within European schools. These guidelines are primarily directed at the relevant policy-makers and at local authorities, secondarily to school-building designers and managers, schoolchildren and their parents, teachers and other school staff.⁹⁸

In Italy, a prevention program for indoor environments is provided in the “Guidelines for the protection and promotion of health in confined environments” (Agreement between the Minister of Health, Regions and Autonomous Provinces - G.U. 27/11/2001).⁹⁹

Moreover, the National Study Group (GdS) on Indoor Air Pollution (established in 2010 at the Italian Institute of Health - ISS) works to provide shared technical–scientific documents in order to allow harmonized actions at national level for improving the correct assessment of indoor air pollution.¹⁰⁰

A perspective from other GARD countries

Exposure to indoor air pollution, typically from multiple sources, is a risk factor for respiratory health, particularly in low resource settings, including people in high income countries in financial austerity. Addressing exposure should improve not only health outcomes, but also the environment, climate and also gender equity, because women typically have larger exposure than men. It will also break the cycle of infants in low resource settings having higher risks of developing CRD in later life. Success depends on developing deep insights into local beliefs and behaviours, and co-creating local primordial, primary and secondary prevention solutions building on the evidence generated about impactful interventions.

Through FRESH AIR and RESPIRE initiatives, data linking indoor air pollution and respiratory health were found and more engaging education about how we breathe and why we get breathless including new short films was created.¹⁰¹ In addition to smoke-related indoor air pollution, the risks of

Table 3 Interventions for CRDs prevention.^{102,105}**Individual level**

- Raising awareness about disease course and risk factors.
- Raising the interaction with the modern health care system.
- Increasing communication with health care providers.
- Raising awareness about usefulness of uninterrupted use of inhalers for maximum benefits.
- Avoiding the use of biomass for cooking or heating and the exposure to ETS.

Community level

- Changing perceptions about CRDs and destigmatizing the disease through educational interventions.
- Raising awareness about the importance of the family in providing physical, economic and psychological support to CRDs patients.

Health system level

- Increasing communication with patients about disease course and risk factors.
- Improving interactions with the community through frontline health care workers.
- Engaging the community with culturally acceptable and locally available media and methods.
- Improving spirometry availability and accessibility in primary care.
- Where spirometry is not available, consider alternatives such as validated questionnaires and tools, particularly in LMICs.
- Implementation of operational primary care guidelines and training programmes for health-care workers and community health workers.
- Improvement of medicine availability and affordability.

Political level

- Investment in strategies to reduce barriers to healthy CRDs behaviours.
- Implementation of enhanced programmes for cleaner fuels and improved appliances.
- Development of CRD-specific health policy with evidence-based protocols, supported by regular availability of and accessibility to medical equipment and medications.
- Define specific indoor air quality management regulations.

Scientific society level

- Continue to increase awareness about CRDs burden and related risk factors.
- Work together to pursue the GARD vision of “a world where all people breathe freely”.

damp and mould on respiratory health have to be considered particularly in times of financial austerity when heating may be reduced and due to the impact of climate change. Researchers in island communities with high humidity and low-lying countries at risk of flooding have been developing innovative examples of community education about the risks and solutions.

Within the RESPIRE collaboration, a study performed in a rural low-literate community in southern India analyzed the lived experiences of people with CRDs: to adopt health behaviours which prevent chronic disease and ensure adequate management, people need to understand the disease, its associated risk factors and treatment options and awareness of the benefits of their health behaviours.¹⁰²

Healthy behaviours are feasible but financial, social or cultural constraints may limit choices. CRDs awareness levels in rural community were limited to symptoms, duration, seasonality and available treatments for symptoms' exacerbation. There was poor understanding of the condition, causes and risk factors for the disease.¹⁰²

People typically were not aware that their disease was not curable or appreciate the efficacy of regular medications; the common practice was to attend nearby medical stores or get some local treatment for temporary relief of symptoms without consulting a doctor.¹⁰²

Biomass fuel was commonly used for cooking; people were unaware that it was a risk factor for their disease and

were happy to continue using it for long-term cultural practice or cost reasons.¹⁰² The use of a more efficient cooking technologies, developed with a participatory approach to meet cooking needs, showed a protective effect on the respiratory health in women and children living in the Central Mexican state of Michoacán.^{103,104}

Improving health literacy and understanding of the disease can lead to early health seeking by the mainstream health care system. The role of HCPs in communicating with patients about risk factors, diagnosis, treatment and disease course is also extremely important. The family plays an important role in providing physical, economic and psychological support and can modulate the patient's health behavior.¹⁰²

Changes in individual knowledge, awareness and attitudes should be accompanied by structural and policy changes (e.g. subsidized tariffs for gas connections, investments to address financial barriers, education and health awareness).¹⁰²

Specific interventions are needed for CRDs prevention at all levels, from patient to health system to policy (Table 3).

Conclusions

In the WHO European Region people spend approximately 90% of their time indoors: 2/3 of this time at home, the rest

at workplace, school and public spaces.¹⁰⁶ Hence, it is important to focus on indoor pollution.

In the absence of specific indoor air quality management regulations, the WHO has developed guidelines for indoor air quality and household fuel combustion to provide health-based recommendations on the best available fuels and technologies for protecting health.^{107–109}

Thus, WHO, scientific societies, patient organizations and other members of the health community should work together to pursue, also for indoor air, the GARD vision of “a world where all people breathe freely” and encourage policy makers to increase their engagement in advocacy for clean air.

Acknowledgements

The authors wish to thank Patrizia Silvi (CNR Institute of Clinical Physiology, Pisa, Italy) for editorial assistance.

This work was supported in part by Ministry of Health in the framework of the National Plan for Complementary Investments (PNC) (project [PREV-A-2022-12377010](https://doi.org/10.1016/j.pulmoe.2021.12.010): MIS-SION).

References

- World Health Organization. Chronic Respiratory Diseases. <https://www.who.int/health-topics/chronic-respiratory-diseases> [Accessed 31 August 2022].
- World Health Organization. Household Air Pollution and Health. <https://www.who.int/news-room/fact-sheets/detail/household-air-pollution-and-health> [Accessed 31 August 2022].
- Health Effects Institute and Institute for Health Metrics and Evaluation. State of Global Air. Health Impacts of Household Air Pollution. <https://www.stateofglobalair.org/health/hap#regional-burden> [Accessed 23 November 2022].
- Viegi G, Taborda-Barata L. A series of narrative reviews on air pollution and respiratory health for Pulmonology: why it is important and who should read it. *Pulmonology*. 2022;28(4):243–4. <https://doi.org/10.1016/j.pulmoe.2021.12.010>.
- De Matteis S, Forastiere F, Baldacci S, Maio S, Tagliaferro S, Fasola S, et al. Issue 1 - “Update on adverse respiratory effects of outdoor air pollution”. Part 1): outdoor air pollution and respiratory diseases: a general update and an Italian perspective. *Pulmonology*. 2022;28(4):284–96. <https://doi.org/10.1016/j.pulmoe.2021.12.008>.
- Sousa AC, Pastorinho MR, Masjedi MR, Urrutia-Pereira M, Arrais M, Nunes E, et al. Issue 1 - “Update on adverse respiratory effects of outdoor air pollution” Part 2): outdoor air pollution and respiratory diseases: perspectives from Angola, Brazil, Canada, Iran, Mozambique and Portugal. *Pulmonology*. 2022;28(5):376–95. <https://doi.org/10.1016/j.pulmoe.2021.12.007>.
- Cavaleiro Rufo J., Annesi-Maesano I., Carreiro-Martins P., Moreira A., Sousa A.C., Pastorinho M.R., et al. Issue 2 - “Update On Adverse Respiratory Effects of Indoor Air pollution”. Part 1): Indoor air Pollution and Respiratory diseases: a General Update and a Portuguese perspective. *Pulmonology*; in press.
- Carrozzi L, Giuliano G, Viegi G, Paoletti P, Di Pede F, Mammini U. The Po River Delta epidemiological study of obstructive lung disease: sampling methods, environmental and population characteristics. *Eur J Epidemiol*. 1990;6(2):191–200. <https://doi.org/10.1007/BF00145793>.
- Baldacci S, Carrozzi L, Viegi G, Giuntini C. Assessment of respiratory effect of air pollution: study design on general population samples. *J Environ Pathol Toxicol Oncol*. 1997;16(2–3):77–83.
- Sapigni T, Biavati P, Simoni M, Viegi G, Baldacci S, Carrozzi L, et al. The Po River Delta Respiratory Epidemiological Survey: an analysis of factors related to level of total serum IgE. *Eur Respirat J*. 1998;11(2):278–83. <https://doi.org/10.1183/09031936.98.11020278>.
- Viegi G, Pedreschi M, Baldacci S, Chiaffi L, Pistelli F, Modena P, et al. Prevalence rates of respiratory symptoms and diseases in general population samples of North and Central Italy. *Int J Tuberc Lung Dis*. 1999;3(11):1034–42.
- Simoni M, Scognamiglio A, Carrozzi L, Baldacci S, Angino A, Pistelli F, et al. Indoor exposures and acute respiratory effects in two general population samples from a rural and an urban area in Italy. *J Expo Anal Environ Epidemiol*. 2004;14(Suppl 1):S144–52. <https://doi.org/10.1038/sj.jea.7500368>.
- Galassi C, De Sario M, Biggeri A, Bisanti L, Chellini E, Ciccone G, et al. Changes in Prevalence of Asthma and Allergies Among Children and Adolescents in Italy: 1994–2002. *Pediatrics*. 2006;117(1):34–42. <https://doi.org/10.1542/peds.2004-2709>.
- Simoni M, Baldacci S, Puntoni R, Pistelli F, Farchi S, Lo Presti E, et al. Respiratory symptoms/diseases and environmental tobacco smoke (ETS) in never smoker Italian women. *Respir Med*. 2007;101(3):531–8. <https://doi.org/10.1016/j.rmed.2006.06.021>.
- Lombardi E, Simoni M, La Grutta S, Viegi G, Bisanti L, Chellini E, et al. Effects of pet exposure in the first year of life on respiratory and allergic symptoms in 7-yr.-old children. The SIDRIA-2 study. *Pediatr Allergy Immunol*. 2010;21(2 Pt 1):268–76. <https://doi.org/10.1111/j.1399-3038.2009.00910.x>.
- Cibella F, Cuttitta G, La Grutta S, Melis MR, Lospalluti ML, Uasuf CG, et al. Proportional Venn diagram and determinants of allergic respiratory diseases in Italian adolescents. *Pediatr Allergy Immunol*. 2011;22(1 Pt 1):60–8. <https://doi.org/10.1111/j.1399-3038.2010.01097.x>.
- Cibella F, Cuttitta G, Della Maggiore R, Ruggieri S, Panunzi S, De Gaetano A, et al. Effect of indoor nitrogen dioxide on lung function in urban environment. *Environ Res*. 2015;138:8–16. <https://doi.org/10.1016/j.envres.2015.01.023>.
- Bentayeb M, Norback D, Bednarek M, Bernard A, Cai G, Cerrai S, et al. Indoor air quality, ventilation and respiratory health in elderly residents living in nursing homes in Europe. *Eur Respir J*. 2015;45(5):1228–38. <https://doi.org/10.1183/09031936.00082414>.
- Ruggieri S, Drago G, Longo V, Colombo P, Balzan M, Bilocca D, et al. Sensitization to dust mite defines different phenotypes of asthma: a multicenter study. *Pediatr Allergy Immunol*. 2017;28(7):675–82. <https://doi.org/10.1111/pai.12768>.
- Drago G, Perrino C, Canepari S, Ruggieri S, L’Abbate L, Longo V, et al. Relationship between domestic smoking and metals and rare earth elements concentration in indoor PM(2.5). *Environ Res*. 2018;165:71–80. <https://doi.org/10.1016/j.envres.2018.03.026>.
- Cilluffo G, Ferrante G, Fasola S, Malizia V, Montalbano L, Ranzi A, et al. Association between asthma control and exposure to greenness and other outdoor and indoor environmental factors: a longitudinal study on a cohort of asthmatic children. *Int J Environ Res Public Health*. 2022;19(1). <https://doi.org/10.3390/ijerph19010512>.
- Viegi G, Carrozzi L, Paoletti P, Vellutini M, Ballerin L, Biavati P, et al. Effects of some indoor environmental factors on respiratory symptoms and lung function in a sample of young non smokers in North Italy. *Aerobiologia (Bologna)*. 1991;7(2):152–9. <https://doi.org/10.1007/BF02270684>.
- Viegi G, Paoletti P, Carrozzi L, Vellutini M, Ballerin L, Biavati P, et al. Effects of home environment on respiratory symptoms and lung function in a general population sample in north Italy. *Eur Respir J*. 1991;4(5):580–6.

24. Viegi G, Carrozzi L, Paoletti P, Vellutini M, DiViggiano E, Baldacci S, et al. Effects of the home environment on respiratory symptoms of a general population sample in middle Italy. *Arch Environ Health*. 1992;47(1):64–70. <https://doi.org/10.1080/00039896.1992.9935946>.
25. Paoletti P, Viegi G, Carrozzi L, Sapigni T, Baldacci S, Pedreschi M, et al. Cigarette smoking, indoor pollution and obstructive lung disease. *Eur Respir Rev*. 1995;5(31):315–22.
26. Simoni M, Carrozzi L, Baldacci S, Pedreschi M, Di Pede F, Angino A, et al. Characteristics of women exposed and unexposed to environmental tobacco smoke (ETS) in a general population sample of North Italy (Po River Delta epidemiological study). *Eur J Epidemiol*. 2001;17(4):363–8. <https://doi.org/10.1023/a:1012797315210>.
27. Simoni M, Carrozzi L, Baldacci S, Scognamiglio A, Di Pede F, Sapigni T, et al. The Po River Delta (north Italy) indoor epidemiological study: effects of pollutant exposure on acute respiratory symptoms and respiratory function in adults. *Arch Environ Health*. 2002;57(2):130–6. <https://doi.org/10.1080/00039890209602928>.
28. Simoni M, Jaakkola MS, Carrozzi L, Baldacci S, Di Pede F, Viegi G. Indoor air pollution and respiratory health in the elderly. *Eur Respir J Suppl*. 2003;40:15s–20s. <https://doi.org/10.1183/09031936.03.00403603>.
29. Simoni M, Lombardi E, Berti G, Rusconi F, La Grutta S, Piffer S, et al. Effects of indoor exposures on respiratory and allergic disorders. *Epidemiol Prev*. 2005;29(2 Suppl):57–61.
30. Simoni M, Lombardi E, Berti G, Rusconi F, La Grutta S, Piffer S, et al. Mould/dampness exposure at home is associated with respiratory disorders in Italian children and adolescents: the SIDRIA-2 Study. *Occup Environ Med*. 2005;62(9):616–22. <https://doi.org/10.1136/oem.2004.018291>.
31. Simoni M, Annesi-Maesano I, Sigsgaard T, et al. School air quality related to dry cough, rhinitis and nasal patency in children. *Eur Respir J*. 2010;35(4):742–9. <https://doi.org/10.1183/09031936.00016309>.
32. Simoni M, Cai G-H, Norback D, Annesi-Maesano I, Lavaud F, Sigsgaard T, et al. Total viable molds and fungal DNA in classrooms and association with respiratory health and pulmonary function of European schoolchildren. *Pediatr Allergy Immunol*. 2011;22(8):843–52. <https://doi.org/10.1111/j.1399-3038.2011.01208.x>.
33. Simoni M, Biavati P, Carrozzi L, Viegi G, Paoletti P, Matteucci G, et al. The Po River Delta (North Italy) indoor epidemiological study: home characteristics, indoor pollutants, and subjects' daily activity pattern. *Indoor Air*. 1998;8(2):70–9. <https://doi.org/10.1111/j.1600-0668.1998.t01-2-00002.x>.
34. Instituto 35. Instituto Nacional de Ecología y Cambio Climático (INECC), <http://airepuro.rowenta.com.mx/principales-contaminantes-del-aire-intramuros/>; 2023 [Accessed 9 January 2023]
35. Salcido GH, Gonzalez-Diaz S, Lira-Quezada CD. Sick building syndrome: how is it affected by airborne pollen, indoor and outdoor pollutants? *Ann Allergy Asthma Immunol*. 2022;129(5, Supplement):S71. <https://doi.org/10.1016/j.anaai.2022.08.706>.
36. Svendsen ER, Gonzales M, Commodore A. The role of the indoor environment: residential determinants of allergy, asthma and pulmonary function in children from a US-Mexico border community. *Sci Total Environ*. 2018;616–617:1513–23. <https://doi.org/10.1016/j.scitotenv.2017.10.162>.
37. Regalado J, Pérez-Padilla R, Sansores R, Páramo Ramirez JI, Brauer M, Paré P, Vedal S. The effect of biomass burning on respiratory symptoms and lung function in rural Mexican women. *Am J Respir Crit Care Med*. 2006;174(8):901–5. <https://doi.org/10.1164/rccm.200503-479OC>.
38. Ramírez-Venegas A, Velázquez-Uncal M, Pérez-Hernández R, NE Guzmán-Bouilloud, Falfán-Valencia R, Mayar-Maya ME, et al. Prevalence of COPD and respiratory symptoms associated with biomass smoke exposure in a suburban area. *Int J Chron Obstruct Pulmon Dis*. 2018;13:1727–34. <https://doi.org/10.2147/COPD.S156409>.
39. Fernández-Plata R, Rojas-Martínez R, Martínez-Briseño D, García-Sancho C, Pérez-Padilla R. Effect of passive smoking on the growth of pulmonary function and respiratory symptoms in schoolchildren. *Rev Invest Clin*. 2016;68(3):119–27.
40. Rivera NYR, Tamayo-Ortiz M, García AM, Just AC, Kloog I, Téllez-Rojo MM, Wright RO, et al. Prenatal and early life exposure to particulate matter, environmental tobacco smoke and respiratory symptoms in Mexican children. *Environ Res*. 2021;192:110365. <https://doi.org/10.1016/j.envres.2020.110365>.
41. Álvarez-Chávez CR, Flores-Bernal JL, Esquer-Peralta J, Munguía-Vega NE, Corella-Madueño MA, Rascón-Careaga A, et al. Detection of allergen sources in the homes of sensitized children. *Environ Health Prev Med*. 2016;21(6):531–8. <https://doi.org/10.1007/s12199-016-0566-7>.
42. Segura-Medina P, Vargas MH, Aguilar-Romero JM, Arreola-Ramírez JL, Miguel-Reyes JL, Salas-Hernández J. Mold burden in house dust and its relationship with asthma control. *Respir Med*. 2019;150:74–80. <https://doi.org/10.1016/j.rmed.2019.02.014>.
43. Gastelum-Arellanez A, Esquivel-Días J, Lopez-Padilla R, Robledo VH, Rodríguez P, Beltran MF, et al. Assessment of persistent indoor VOCs inside public transport during winter season. *Chemosphere*. 2021;263:128127. <https://doi.org/10.1016/j.chemosphere.2020.128127>.
44. Baggio G, Tejada CAO, dos Santos AMA, Triaca LM. Indoor air pollution and respiratory diseases in rural areas of North and Northeast Brazil. *Int J Econ Finance*. 2019;11:66–75.
45. Ponte EV, Lima A, Almeida PCA, de Jesus JPV, Souza-Machado A, Barreto ML, et al. Rural to urban migration contributes to the high burden of asthma in the urban area. *Clin Respir J*. 2019;13(9):560–6. <https://doi.org/10.1111/crj.13058>.
46. Fernandes AGO, de Souza-Machado C, Pinheiro GP, de Oliva ST, Mota RCL, de Lima VB, et al. Dual exposure to smoking and household air pollution is associated with an increased risk of severe asthma in adults in Brazil. *Clin Transl Allergy*. 2018;8:48. <https://doi.org/10.1186/s13601-018-0235-6>.
47. Hystad P, Duong M, Brauer M, Larkin A, Arku R, Kurmi OP, et al. Health effects of household solid fuel use: findings from 11 countries within the prospective urban and rural epidemiology study. *Environ Health Perspect*. 2019;127(5):57003. <https://doi.org/10.1289/EHP3915>.
48. da Silva LFF, Saldiva SRDM, Saldiva PHN, Dolhnikoff M. Impaired lung function in individuals chronically exposed to biomass combustion. *Environ Res*. 2012;112:111–7. <https://doi.org/10.1016/j.envres.2011.10.012>.
49. Gioda A, Tonietto GB, Leon AP. Exposure to the use of firewood for cooking in Brazil and its relation with the health problems of the population. *Cien Saude Colet*. 2019;24(8):3079–88. <https://doi.org/10.1590/1413-81232018248.23492017>.
50. Galduróz JC, Tomita N, Bezerra AG. Measures to reduce smoking: Brazil takes the lead. *Braz J Psychiatry*. 2020;42(4):456–7. <https://doi.org/10.1590/1516-4446-2020-0853>.
51. Schivinski CIS, de Assumpção MS, de Figueiredo FCXS, Wamosy RMG, Ferreira LG, Ribeiro JD. Impulse oscillometry, spirometry, and passive smoking in healthy children and adolescents. *Rev Port Pneumol*. 2006;23(6):311–6. <https://doi.org/10.1016/j.rppnen.2017.06.005>. 2017.
52. Freire APC, Ramos D, Leite MR, Silva BS, David RM, Uzeloto JS, et al. Influence of time and frequency of passive smoking exposure on mucociliary clearance and the autonomic nervous system. *Respir Care*. 2016;61(4):453–61. <https://doi.org/10.4187/respcare.04398>.
53. Schvartsman C, Farhat SCL, Schvartsman S, Saldiva PHN. Parental smoking patterns and their association with wheezing in children. *Clinics*. 2013;68(7):934–9. [https://doi.org/10.6061/clinics/2013\(07\)08](https://doi.org/10.6061/clinics/2013(07)08).

54. SC Fernandes S de, Solé D, Camargos P, de Andrade CR, Ibiapina C da C. Factors associated with asthma expression in adolescents. *J Bras Pneumol.* 2018;44(1):12–7. <https://doi.org/10.1590/S1806-37562017000000078>.
55. Neto ACP, Solé D, Hirakata V, Schmid LS, Klock C, Barreto SSM. Risk factors for asthma in schoolchildren in Southern Brazil. *Allergol Immunopathol (Madr)*. 2020;48(3):237–43. <https://doi.org/10.1016/j.aller.2019.07.003>.
56. Sousa NA, Segalin B, Busse AL, Filho WJ, Fornaro A, Gonçalves FLT. Indoor/outdoor particulate matter and health risk in a nursing community home in São Paulo, Brazil. *Atmosph Pollut Res.* 2021;12(10):101188. <https://doi.org/10.1016/j.apr.2021.10.1188>.
57. Segalin B, Kumar P, Micadei K, Fornaro A, Gonçalves FLT. Size-segregated particulate matter inside residences of elderly in the Metropolitan Area of São Paulo, Brazil. *Atmosph Environ.* 2017;148:139–51. <https://doi.org/10.1016/j.atmosenv.2016.10.004>.
58. Huy LN, Oanh NTK, Phuc NH, Nhung CP. Survey-based inventory for atmospheric emissions from residential combustion in Vietnam. *Environ Sci Pollut Res.* 2021;28(9):10678–95. <https://doi.org/10.1007/s11356-020-11067-6>.
59. Van Minh H, Giang KB, Ngoc NB, et al. Prevalence of tobacco smoking in Vietnam: findings from the Global Adult Tobacco Survey 2015. *Int J Public Health.* 2017;62(Suppl 1):121–9. <https://doi.org/10.1007/s00038-017-0955-8>.
60. Quang TN. Fine and ultrafine particle concentrations at Hanoi primary schools. *J Sci Technol Civil Eng.* 2017;11(6):162–7.
61. Quang TN, Hue NT, Thai P, Mazaheri M, Morawska L. Exploratory assessment of indoor and outdoor particle number concentrations in Hanoi households. *Sci Total Environ.* 2017;599–600:284–90. <https://doi.org/10.1016/j.scitotenv.2017.04.154>.
62. Lâm HT, Ekerljung L, Bjerg A, Vän T Tng N, Lundbäck B, Rönmark E. Sensitization to airborne allergens among adults and its impact on allergic symptoms: a population survey in northern Vietnam. *Clin Transl Allergy.* 2014;4(1):6. <https://doi.org/10.1186/2045-7022-4-6>.
63. Sy D, Thanh Binh M, Quoc N, Hung NV, Quynh Nhu DT, Bao NQ, et al. Prevalence of asthma and asthma-like symptoms in Dalat Highlands, Vietnam. *Singapore Med J.* 2007;48(4):294–303.
64. Matsui EC, Perzanowski M, Peng RD, et al. Effect of an integrated pest management intervention on asthma symptoms among mouse-sensitized children and adolescents with asthma: a randomized clinical trial. *JAMA.* 2017;317(10):1027–36. <https://doi.org/10.1001/jama.2016.21048>.
65. Cần Trọng Với ô Nhiễm Không Khí Trong Nhà. <https://nld.com.vn/news-2020108220732842.htm> [Accessed 15 March 2023].
66. Johnson P, Balakrishnan K, Ramaswamy P, Ghosh S, Sadhasivam M, Abirami O, et al. Prevalence of chronic obstructive pulmonary disease in rural women of Tamilnadu: implications for refining disease burden assessments attributable to household biomass combustion. *Glob Health Action.* 2011;4:7226. <https://doi.org/10.3402/gha.v4i0.7226>.
67. Agrawal S. Effect of indoor air pollution from biomass and solid fuel combustion on prevalence of self-reported asthma among adult men and women in India: findings from a nationwide large-scale cross-sectional survey. *J Asthma.* 2012;49(4):355–65. <https://doi.org/10.3109/02770903.2012.663030>.
68. Rao SR, Chitra GA, Elavarasu G, Kamaraj P, Kaliaperumal K, Kaur P. Exposure to mosquito coil and biomass fuel smoke and respiratory health in rural Tamil Nadu, India. *J Public Health (Oxf)*. 2022;44(3):625–33. <https://doi.org/10.1093/pubmed/fdab119>.
69. Bhat RY, Manjunath N. Correlates of acute lower respiratory tract infections in children under 5 years of age in India. *Int J Tuberc Lung Dis.* 2013;17(3):418–22. <https://doi.org/10.5588/ijtld.12.0117>.
70. Naz S, Page A, Agho KE. Potential impacts of modifiable behavioral and environmental exposures on reducing burden of under-five mortality associated with household air pollution in Nepal. *Matern Child Health J.* 2018;22(1):59–70. <https://doi.org/10.1007/s10995-017-2355-3>.
71. Acharya P, Mishra SR, Berg-Beckhoff G. Solid fuel in kitchen and acute respiratory tract infection among under five children: evidence from Nepal demographic and health survey 2011. *J Community Health.* 2015;40(3):515–21. <https://doi.org/10.1007/s10900-014-9965-0>.
72. Kurmi OP, Devereux GS, Smith WCS, Semple S, Steiner MF, Simkhada P, et al. Reduced lung function due to biomass smoke exposure in young adults in rural Nepal. *Eur Respir J.* 2013;41(1):25–30. <https://doi.org/10.1183/09031936.00220511>.
73. Sinha B, Singla R, Chowdhury R. An epidemiological profile of chronic obstructive pulmonary disease: a community-based study in Delhi. *J Postgrad Med.* 2017;63(1):29–35. <https://doi.org/10.4103/0022-3859.194200>.
74. Deka H, Mahanta P, Ahmed SJ, Rajbangshi MC, Konwar R, Basumatari B. Risk factors of childhood asthma among patients attending a tertiary care centre in North-East India. *J Asthma Allergy.* 2022;15:1293–303. <https://doi.org/10.2147/JAA.S374007>.
75. Gupta D, Aggarwal AN, Chaudhry K, Chhabra SK, D'Souza GA, Jindal SK, et al. Household environmental tobacco smoke exposure, respiratory symptoms and asthma in non-smoker adults: a multicentric population study from India. *Indian J Chest Dis Allied Sci.* 2006;48(1):31–6.
76. Melsom T, Brinch L, Hessen JO, Schei MA, Kolstrup N, Jacobsen BK, et al. Asthma and indoor environment in Nepal. *Thorax.* 2001;56(6):477–81. <https://doi.org/10.1136/thorax.56.6.477>.
77. Dahal GP, Johnson FA, Padmadas SS. Maternal smoking and acute respiratory infection symptoms among young children in Nepal: multilevel analysis. *J Biosoc Sci.* 2009;41(6):747–61. <https://doi.org/10.1017/S002193200990113>.
78. Thapa P, Pandey AR, Dhungana RR, Bista B, Thapa B, Mishra SR. Risk of ARI among non-exclusively breastfed under-five passive smoker children: a hospital-based cross-sectional study of Nepal. *Front Public Health.* 2016;4:23. <https://doi.org/10.3389/fpubh.2016.00023>.
79. Sooronbaev T, Ghezai B, Williams S, Almatalieva M, Mademilov M, Chavannes N. Indoor pollutions and the prevalence of respiratory symptoms among highlanders in Kyrgyzstan. *Eur Respir J.* 2015;46(suppl 59). <https://doi.org/10.1183/13993003.congress-2015.PA4094>.
80. Tabyshova A, Emilov B, Seiitov M, Beyshenbekova A, Osmonov B, Estebesova B, et al. Indoor air pollution and respiratory symptoms, and pneumonia in highland children in Kyrgyzstan. *Eur Respir J.* 2017;50(suppl 61). <https://doi.org/10.1183/1393003.congress-2017.PA1859>.
81. Brakema EA, Tabyshova A, Kasteleyn MJ, Molendijk E, van der Kleij RMJJ, van Boven JFM, et al. High COPD prevalence at high altitude: does household air pollution play a role? *Eur Respir J.* 2019;53(2). <https://doi.org/10.1183/13993003.01193-2018>.
82. Tabyshova A, Estebesova B, Beishenbekova A, Sooronbaev T, Brakema EA, Chavannes NH, et al. Clinical Characteristics, Treatment Patterns and Economic Burden of COPD in Kyrgyzstan: a FRESH AIR Study. *Int J Chron Obstruct Pulmon Dis.* 2021;16:2833–43. <https://doi.org/10.2147/COPD.S322778>.
83. van Gemert F, de Jong C, Kirenga B, Musinguzi P, Buteme S, Sooronbaev T, et al. Effects and acceptability of implementing improved cookstoves and heaters to reduce household air pollution: a FRESH AIR study. *NPJ Prim Care Respir Med.* 2019;29(1):32. <https://doi.org/10.1038/s41533-019-0144-8>.
84. World Health Organization. Evidence, Policy, Impact: WHO Guide For Evidence-Informed Decision-Making. <https://apps.who.int/iris/handle/10665/350994> [Accessed 16 March 2023].

85. International Primary Care Respiratory Group. FRESH AIR. <https://www.ipcr.org/freshair/> [Accessed 22 February 2023].
86. van Gemert F, Kirenga B, Chavannes N, Kamya M, Luzige S, Musinguzi P, et al. Prevalence of chronic obstructive pulmonary disease and associated risk factors in Uganda (FRESH AIR Uganda): a prospective cross-sectional observational study. *Lancet Glob Health*. 2015;3(1):e44–51. [https://doi.org/10.1016/S2214-109X\(14\)70337-7](https://doi.org/10.1016/S2214-109X(14)70337-7).
87. Adeloye D, Song P, Zhu Y, Campbell H, Sheikh A, Rudan I. Global, regional, and national prevalence of, and risk factors for, chronic obstructive pulmonary disease (COPD) in 2019: a systematic review and modelling analysis. *Lancet Respir Med*. 2022;10(5):447–58. [https://doi.org/10.1016/S2213-2600\(21\)00511-7](https://doi.org/10.1016/S2213-2600(21)00511-7).
88. Hanafi NS, Agarwal D, Chippagiri S, Brakema EA, Pinnock H, Sheikh A, et al. Chronic respiratory disease surveys in adults in low- and middle-income countries: a systematic scoping review of methodological approaches and outcomes. *J Glob Health*. 2021;11:04026. <https://doi.org/10.7189/jogh.11.04026>.
89. Cragg L, Williams S, Chavannes NH. FRESH AIR: an implementation research project funded through Horizon 2020 exploring the prevention, diagnosis and treatment of chronic respiratory diseases in low-resource settings. *NPJ Prim Care Respir Med*. 2016;26:16035. <https://doi.org/10.1038/nnpjcrm.2016.35>.
90. Brakema EA, van der Kleij RMJJ, Poot CC, An PL, Anastasaki M, Crone MR, et al. Mapping low-resource contexts to prepare for lung health interventions in four countries (FRESH AIR): a mixed-method study. *Lancet Global Health*. 2022;10(1):e63–76. [https://doi.org/10.1016/S2214-109X\(21\)00456-3](https://doi.org/10.1016/S2214-109X(21)00456-3).
91. Brakema EA, van Gemert FA, Williams S, Sooronbaev T, Emilov B, Mademilov M, et al. Implementing a context-driven awareness programme addressing household air pollution and tobacco: a FRESH AIR study. *NPJ Prim Care Respir Med*. 2020;30(1):42. <https://doi.org/10.1038/s41533-020-00201-z>.
92. Anastasaki M, Tsiligianni I, Sifaki-Pistola D, Chatzea VE, Karelis A, Bertias A, et al. Household air pollution and respiratory health in rural crete, greece: a cross-sectional FRESH AIR study. *Atmosphere (Basel)*. 2021;12(11). <https://doi.org/10.3390/atmos12111369>.
93. International Primary Care Respiratory Group. GHRN Publications. <https://www.ipcr.org/ghrn-publications/> [Accessed 22 February 2023].
94. Boudewijns EA, Trucchi M, van der Kleij RMJJ, Vermond D, Hoffman CM, Chavannes NH, et al. Facilitators and barriers to the implementation of improved solid fuel cookstoves and clean fuels in low-income and middle-income countries: an umbrella review. *Lancet Planetary Health*. 2022;6(7):e601–12. [https://doi.org/10.1016/S2542-5196\(22\)00094-8](https://doi.org/10.1016/S2542-5196(22)00094-8).
95. McCarron A, Uny I, Caes L, Lucas SE, Semple S, Ardrey J, et al. Solid fuel users' perceptions of household solid fuel use in low- and middle-income countries: a scoping review. *Environ Int*. 2020;143:105991. <https://doi.org/10.1016/j.envint.2020.105991>.
96. Ministero della Salute. Documento Di Strategia GARD-I. <https://www.salute.gov.it/portale/gard/dettaglioPubblicazioniGard.jsp?id=1206&lingua=italiano/> [Accessed 16 March 2023].
97. Ministero della Salute. La qualità dell'aria nelle scuole e rischi per malattie respiratorie e allergiche. Quadro conoscitivo sulla situazione italiana e strategie di prevenzione, https://www.salute.gov.it/portale/documentazione/p6_2_2_1.jsp?id=1892&lingua=italiano; 2023 [Accessed 16 March 2023].
98. Kephelopoulou S, Csobod E, Bruinen DBY, De OFE. Guidelines for healthy environments within European schools. JRC Public Reposit. 2014. <https://doi.org/10.2788/89936>.
99. Gazzetta ufficiale della Repubblica Italiana. Accordo tra il Ministro della salute, le regioni e le province autonome sul documento concernente: “Linee-guida per la tutela e la promozione della salute negli ambienti confinati”, <https://www.gazzettaufficiale.it/eli/id/2001/11/27/01A11196/sg/>; 2023 [accessed March 16, 2023].
100. Settimo G, Manigrasso M, Avino P. Indoor air quality: a focus on the european legislation and state-of-the-art research in Italy. *Atmosphere*. 2020;11(4). <https://doi.org/10.3390/atmos11040370>.
101. International Primary Care Respiratory Group. How We Breathe. <https://www.ipcr.org/howwebreathe/> [Accessed 22 February 2023].
102. Paul B, Isaac R, Hemalatha R, Jebaraj P, Muthathal S, Das D, et al. Development of an educational intervention to reduce the burden of adult chronic lung disease in rural India: inputs from a qualitative study. *PLoS ONE*. 2021;16(7):e0254534. <https://doi.org/10.1371/journal.pone.0254534>.
103. Romieu I, Riojas-Rodríguez H, Marrón-Mares AT, Schilman A, Perez-Padilla R, Masera O. Improved biomass stove intervention in rural Mexico: impact on the respiratory health of women. *Am J Respir Crit Care Med*. 2009;180(7):649–56. <https://doi.org/10.1164/rccm.200810-1556OC>.
104. Schilman A, Riojas-Rodríguez H, Ramírez-Sedeño K, Berrueta VM, Pérez-Padilla R, Romieu I. Children's respiratory health after an efficient biomass stove (Patsari) intervention. *Ecohealth*. 2015;12(1):68–76. <https://doi.org/10.1007/s10393-014-0965-4>.
105. Rossaki FM, Hurst JR, van Gemert F, Kirenga BJ, Williams S, Khoo EM, et al. Strategies for the prevention, diagnosis and treatment of COPD in low-and middle- income countries: the importance of primary care. *Expert Rev Respir Med*. 2021;15(12):1563–77. <https://doi.org/10.1080/17476348.2021.1985762>.
106. World Health Organization. Regional Office for Europe. Combined or Multiple Exposure to Health Stressors in Indoor Built Environments: An Evidence-Based Review Prepared for the WHO Training Workshop “Multiple Environmental Exposures and Risks. Bonn, Germany: World Health Organization. Regional Office for Europe; 2014. 16–18 October 2013 <https://apps.who.int/iris/handle/10665/350495/> [Accessed 16 March 2023].
107. World Health Organization. WHO Guidelines for Indoor Air Quality: Household Fuel Combustion. <https://apps.who.int/iris/handle/10665/141496/> [Accessed 16 March 2023].
108. World Health Organization. WHO Guidelines for Indoor Air Quality: Selected Pollutants. <https://www.who.int/publications/i/item/9789289002134/> [Accessed 16 March 2023].
109. World Health Organization. WHO Guidelines for Indoor Air Quality: Dampness and Mould. <https://www.who.int/publications/i/item/9789289041683/> [Accessed 16 March 2023].