



## EDITORIAL

## Combination of cryobiopsy with EBUS-TBNA—Might rapid on-site evaluation successfully drive patient selection?



Endosonography, which is a minimally invasive technique adopted for the diagnosis of hilar and mediastinal lymphadenopathies, is the gold standard for mediastinal staging of non-small cell lung cancer (NSCLC).<sup>1,2</sup>

Small cytological samples collected by needle-based endosonography may be inadequate to describe histopathology of lymphoproliferative disorders and to assess molecular patterns in patients with advanced lung cancer.<sup>3–5</sup>

Needles of different size and alternative sampling modalities, such as endobronchial ultrasound-guided intranodal forceps biopsy (EBUS-IFB), have been proposed to overcome this limitation<sup>2,6,7</sup>; however, studies have shown that different needle size<sup>5</sup> do not affect overall diagnostic yield and tissue sampling.<sup>2,6,7</sup> The addition of EBUS-IFB to endobronchial ultrasound-guided transbronchial needle aspiration (EBUS-TBNA) may increase the diagnostic yield in cases of sarcoidosis and lymphoma.<sup>8</sup> Unfortunately, the complication rate of the combined approach is higher when compared with EBUS-TBNA alone.<sup>8</sup>

Mediastinal cryobiopsy has emerged as a promising technology to improve the diagnostic yield of mediastinal lymphadenopathy.<sup>9–12</sup>

Experimental and observational studies which compared EBUS-cryobiopsy with EBUS-TBNA demonstrated the added diagnostic value of cryobiopsy, without any increased rate of adverse events<sup>9–12</sup>; however, no differences in the accuracy of the diagnosis of lung cancer were detected. Cryobiopsy was associated with a higher sensitivity in the diagnosis of uncommon malignancies, lymphoproliferative disorders, and benign lesions; moreover, the material collected with cryobiopsy in cases of advanced NSCLC was more suitable for molecular analysis.<sup>9–12</sup>

In a recent issue of the Journal, two studies evaluated the utility of EBUS-cryobiopsy in combination with EBUS-TBNA.<sup>13,14</sup>

Cheng et al. found that both cryobiopsy and nodal forceps biopsy can increase the diagnostic yield for mediastinal lesions when added to EBUS-TBNA.<sup>13</sup> No differences were found for lung cancers but the sensitivity was higher in the

diagnosis of uncommon tumours.<sup>13</sup> The inclusion of cryobiopsy in the diagnostic process resulted in a higher yield for benign diseases and improved the collection of biological materials for molecular testing in those with NSCLC. Few minor adverse events were recorded.<sup>13</sup>

Poletti et al. retrospectively reviewed 48 patients who underwent EBUS-cryobiopsy after EBUS-TBNA.<sup>14</sup> The Authors described a higher yield of cryobiopsy (95.8 %) but the reported sensitivity of EBUS-TBNA (54 %) was lower than previously described.<sup>9–14</sup> Notably, the diagnosis of hyperplastic lymphadenopathy was considered formally acceptable only when specific morphologic criteria based on the recognition of lymph node architecture were identifiable.

No differences were detected in the diagnosis of lung cancer, but a higher yield of cryobiopsy was demonstrated in sarcoidosis, lymphoproliferative disorders, rare mediastinal disorders and hyperplastic lymphadenopathies. Furthermore, pathologists favoured cryobiopsy samples for molecular analysis in patients with advanced NSCLC. No adverse events were reported.<sup>14</sup>

The studies confirmed previous findings on the safety and accuracy of the technique showing the advantages of cryoprobe in comparison with forceps biopsy: its added value can be proved when needle aspiration fails to provide enough material for molecular analysis of NSCLC and when a definitive diagnosis of benign diseases and less common mediastinal tumours has to be confirmed.<sup>13,14</sup>

However, some critical points need to be clarified, from the procedures to patients who may really benefit.<sup>15,16</sup>

Cheng et al. performed EBUS-TBNA (four passes), then cryobiopsy (one sampling) and forceps biopsy (3 samplings) using a randomized approach.<sup>13</sup> An incision in the tracheobronchial wall was performed with an electro-surgical knife to insert cryoprobe and forceps into the mediastinum during spontaneous breathing, with conscious sedation and without intubation (oral/nasal route).<sup>13</sup> A complicated procedure including the use of an electro-surgical knife, not available in the majority of the centers, was carried out by expert bronchoscopists.<sup>13</sup>

<https://doi.org/10.1016/j.pulmoe.2024.02.008>

2531-0437/© 2024 Sociedade Portuguesa de Pneumologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Poletti et al. described an easier technique passing cryoprobe through the puncture site of EBUS-TBNA. All the procedures were performed using rigid bronchoscopes. The number of samplings was not reported.<sup>14</sup>

No studies have compared diagnostic yield, safety profile, and patient comfort during cryo-EBUS using different sedation protocols and endoscope insertion routes. Different strategies of nodal cryo-sampling have been described in the literature.<sup>11–14,17</sup> However, the best technique in terms of probe insertion, adequate number of biopsies, and cooling time has not been determined yet.

A standardized, easy-to-perform procedure would be needed to achieve a widespread adoption.

In both studies cryobiopsy was added to EBUS-TBNA and the use of rapid on-site evaluation (ROSE)<sup>2,18</sup> was not reported.

The need for an additional sampling tool (i.e., cryo- or forceps biopsy) might be superfluous if ROSE is included in the diagnostic workflow of endosonography: ROSE does not increase the diagnostic yield<sup>2,18,19</sup> but can provide immediate information on the quantity and quality of the material. It can reduce the number of needle passes and can improve sample adequacy for diagnosis and molecular profiling or might guide subsequent sampling with cryoprobe in case of poor collection, suspected uncommon tumours or lymphoproliferative disorders.<sup>2,18,19</sup>

In conclusion, the studies by Cheng et al. and Poletti et al. highlight the potential utility of mediastinal cryobiopsy in comparison with forceps biopsy as a complementary tool to overcome limitations of conventional EBUS-TBNA.<sup>13,14</sup>

Future studies including ROSE could help understand the subgroups of patients with mediastinal lymphadenopathies who can have a diagnostic advantage from cryobiopsy. Larger studies should assess the accuracy of this technique in the diagnosis of lymphomas and benign lesions. The role of mediastinal cryobiopsy to identify genomic background in common epithelial tumours represents another interesting field of investigation.

Standardization of the technique is needed before its wider adoption in daily clinical practice.

## Authorship

M.M., M.W. and G.S.: Conception and design; drafting the article; final approval of the version to be submitted.

## Conflicts of interest

None of the Authors has conflict of interest to disclose.

## Funding

None.

## References

- Vilman P, Clementsen PF, Colella S, Siemsen M, De Leyn P, Dumonceau JM, et al. Combined endobronchial and oesophageal endosonography for the diagnosis and staging of lung cancer. *European Society of Gastrointestinal Endoscopy (ESGE) Guideline, in cooperation with the European Respiratory Society (ERS) and the European Society of Thoracic Surgeons (ESTS)*. *Eur Respir J*. 2015;46(1):40–60. <https://doi.org/10.1183/09031936.00064515>.
- Wahidi MM, Herth F, Yasufuku K, Shepherd RW, Yarmus L, Chawla M, et al. Technical aspects of endobronchial ultrasound-guided transbronchial needle aspiration: CHEST guideline and expert panel report. *Chest*. 2016;149(3):816–35. <https://doi.org/10.1378/chest.15-1216>.
- Labarca G, Sierra-Ruiz M, Kheir F, Folch E, Majid A, Mehta HJ, et al. Diagnostic accuracy of endobronchial ultrasound transbronchial needle aspiration in lymphoma. A systematic review and meta-analysis. *Ann Am Thorac Soc*. 2019;16(11):1432–9. <https://doi.org/10.1513/AnnalsATS.201902-1750C>.
- Labarca G, Folch E, Jantz M, Mehta HJ, Majid A, Fernandez-Bussy S. Adequacy of samples obtained by endobronchial ultrasound with transbronchial needle aspiration for molecular analysis in patients with non-small cell lung cancer. Systematic review and meta-analysis. *Ann Am Thorac Soc*. 2018;15(10):1205–16. <https://doi.org/10.1513/AnnalsATS.201801-0450C>.
- Zhao JJ, Chan HP, Soon YY, Huang Y, Soo RA, Kee ACL. A systematic review and meta-analysis of the adequacy of endobronchial ultrasound transbronchial needle aspiration for next-generation sequencing in patients with non-small cell lung cancer. *Lung Cancer*. 2022;166:17–26. <https://doi.org/10.1016/j.lungcan.2022.01.018>.
- Wolters C, Darwiche K, Franzen D, Hager T, Bode-Lesnievska B, Kneuert PJ, et al. A prospective, randomized trial for the comparison of 19-G and 22-G endobronchial ultrasound-guided transbronchial aspiration needles; introducing a novel end point of sample weight corrected for blood content. *Clin Lung Cancer*. 2019;20(3):e265–73. <https://doi.org/10.1016/j.clcc.2019.02.019>.
- Sood R, Alape D, Thakkar D, Shadchehr S, Acash G, Tronic BJ, et al. Comparison of sample adequacy and diagnostic yield of the 21-G and 25-G EBUS TBNA needles. *J Bronchology Interv Pulmonol*. 2022;29(1):34–8. <https://doi.org/10.1097/LBR.0000000000000753>.
- Agrawal A, Ghori U, Chaddha U, Murgu S. Combined EBUS-IFB and EBUS-TBNA vs EBUS-TBNA alone for intrathoracic adenopathy: a meta-analysis. *Ann Thorac Surg*. 2022;114(1):340–8. <https://doi.org/10.1016/j.athoracsur.2020.12.049>.
- Botana-Rial M, Lojo-Rodríguez I, Leiro-Fernández V, Ramos-Hernández C, González-Montaos A, Pazos-Area L, et al. Is the diagnostic yield of mediastinal lymph node cryobiopsy (cryoEBUS) better for diagnosing mediastinal node involvement compared to endobronchial ultrasound-guided transbronchial needle aspiration (EBUS-TBNA)? A systematic review. *Respir Med*. 2023;218:107389. <https://doi.org/10.1016/j.rmed.2023.107389>.
- Ariza-Protá M, Pérez-Pallarés J, Fernández-Fernández A, García-Alfonso L, Cascón JA, Torres-Rivas H, et al. Endobronchial ultrasound-guided transbronchial mediastinal cryobiopsy in the diagnosis of mediastinal lesions: safety, feasibility and diagnostic yield - experience in 50 cases. *ERJ Open Res*. 2023;9(2):00448–2022. <https://doi.org/10.1183/23120541.00448-2022>.
- Zhang J, Guo JR, Huang ZS, Fu WL, Wu XL, Wu N, et al. Transbronchial mediastinal cryobiopsy in the diagnosis of mediastinal lesions: a randomised trial. *Eur Respir J*. 2021;58(6):2100055. <https://doi.org/10.1183/13993003.00055-2021>.
- Fan Y, Zhang AM, Wu XL, Huang ZS, Kontogianni K, Sun K, et al. Transbronchial needle aspiration combined with cryobiopsy in the diagnosis of mediastinal diseases: a multicentre, open-label, randomised trial. *Lancet Respir Med*. 2023;11(3):256–64. [https://doi.org/10.1016/S2213-2600\(22\)00392-7](https://doi.org/10.1016/S2213-2600(22)00392-7).

13. Cheng TL, Huang ZS, Zhang J, Wang J, Zhao J, Kontogianni K, et al. Comparison of cryobiopsy and forceps biopsy for the diagnosis of mediastinal lesions: a randomised clinical trial. *Pulmonology*. 2024;S2531-0437(23):00240–4. <https://doi.org/10.1016/j.pulmoe.2023.12.002>.
14. Poletti V, Petrarulo S, Piciucchi S, Dubini A, De Grauw AJ, Sultani F, et al. EBUS-guided cryobiopsy in the diagnosis of thoracic disorders. *Pulmonology*. 2024;S2531-0437(23):00223–4. <https://doi.org/10.1016/j.pulmoe.2023.11.008>.
15. Mondoni M, Sotgiu G. Optimizing the endoscopic diagnosis of mediastinal lymphadenopathy: a glimpse on cryobiopsy. *BMC Pulm Med*. 2022;22(1):355. <https://doi.org/10.1186/s12890-022-02160-2>.
16. Annema JT, Rozman A. Endobronchial ultrasound-guided cryobiopsy-when is it indicated? *Lancet Respir Med*. 2023;11(3):e23. [https://doi.org/10.1016/S2213-2600\(22\)00486-6](https://doi.org/10.1016/S2213-2600(22)00486-6).
17. Ariza-Prota MA, Pérez-Pallarés J, Fernández-Fernández A, López-González F, Cascón JA, García-Alfonso L, et al. Transbronchial mediastinal cryobiopsy in the diagnosis of mediastinal lymph nodes: a case series - how to do it. *Arch Bronconeumol*. 2022;58(10):718–21. <https://doi.org/10.1016/j.arbres.2022.05.006>.
18. Sehgal IS, Dhooria S, Aggarwal AN, Agarwal R. Impact of Rapid On-Site Cytological Evaluation (ROSE) on the diagnostic yield of transbronchial needle aspiration during mediastinal lymph node sampling: systematic review and meta-analysis. *Chest*. 2018;153(4):929–38. <https://doi.org/10.1016/j.chest.2017.11.004>.
19. Mondoni M, Gasparini S, Varone F, Trisolini R, Mancino L, Rossi G, et al. Accuracy and predictors of success of EUS-B-FNA in the diagnosis of pulmonary malignant lesions: a prospective multi-center Italian study. *Respiration*. 2022;101(8):775–83. <https://doi.org/10.1159/000524398>.

M. Mondoni<sup>a,\*</sup>, M.M. Wahidi<sup>b</sup>, G. Sotgiu<sup>c</sup>

<sup>a</sup> *Respiratory Unit, ASST Santi Paolo e Carlo, Department of Health Sciences, Università degli Studi di Milano, Milan, Italy*

<sup>b</sup> *Northwestern University Feinberg School of Medicine, Chicago, IL, USA*

<sup>c</sup> *Clinical Epidemiology and Medical Statistics Unit, Department of Medical, Surgical and Experimental Sciences, University of Sassari, Sassari, Italy*

\* Corresponding author at: Respiratory Unit, ASST Santi Paolo e Carlo, Department of Health Sciences, Università degli Studi di Milano, Via A. Di Rudinì n.8, 20142, Milan, Italy. E-mail address: [michele.mondoni@asst-santipaolocarlo.it](mailto:michele.mondoni@asst-santipaolocarlo.it) (M. Mondoni).

Received 12 February 2024; Accepted 15 February 2024  
Available online 27 March 2024