

Although it can be expelled spontaneously,⁷ an invasive method may be required when the needle cannot be reached by bronchoscopy. Otherwise, the needle may perforate or migrate to the vascular system or cause chronic inflammation as a response to a foreign object. Only in 7 of the 22 cases reported did the needle break after the adenopathy was punctured [locations 7 (3 patients), 4L (2), 4R (1) and 10R (1)]. The potential causes of needle breakage include⁷: (1) tangential position of the needle due to the angle required for puncture; (2) harder than usual cartilaginous rings or lymph nodes; (3) manufacturing defects of the needle. To prevent this complication, excessive angulation of the needle should be avoided and its structural integrity should be checked prior to each puncture. The needle can also break and remain retained within the sheath.⁸ In this case, when the sheath is extracted from the working channel and the stiffening wire is inserted, the broken distal end of the needle will be expelled. Needles generally break prior to the procedure.

The lesson learned from this experience is that, although EBUS is a safe technique, complications – such as transbronchial puncture needle breakage – may arise. The structural integrity of the needles should be checked throughout the procedure and upon completion. The needle must be extracted. If it cannot be recovered by conventional bronchofibroscopy, an invasive method will be required.

Abbreviations

EBUS	endobronchial ultrasound
EBUS-TBNA	endobronchial ultrasound with real-time guided transbronchial needle aspiration

Authors' contributions

VR, AG and LV were responsible for the conception and design of the study, and wrote and edited the manuscript. AC contributed to the drafting and revision of the manuscript. All authors read and approved the final manuscript.

Funding

This study was undertaken without funding.

Conflicts of interest

The author has no conflicts of interest to declare.

References

1. Detterbeck FC, Boffa DJ, Kim AW, Tanoue LT. The Eighth Edition Lung Cancer Stage Classification. *Chest*. 2017;151:193–203.
2. Eapen GA, Shah AM, Lei X, Jimenez CA, Morice RC, Yarmus L, et al. Complications, consequences, and practice patterns of endobronchial ultrasound-guided transbronchial needle aspiration. Results of the AQUIRE registry. *Chest*. 2013;143:1044–53.
3. Vaidya PJ, Munavvar M, Leuppi JD, Mehta AC, Chhajed PN. Endobronchial ultrasound-guided transbronchial needle aspiration: safe as it sounds. *Respirology*. 2017;22:1093–101.
4. Asano F, Aoe M, Ohsaki Y, Okada Y, Sasada S, Sato S, et al. Complications associated with endobronchial ultrasound-guided transbronchial needle aspiration: a nationwide survey by the Japan Society for Respiratory Endoscopy. *Respir. Res.* 2013;14:50.
5. Özgüll MA, Çetinkaya E, Tutar N, Özgüll G. An unusual complication of endobronchial ultrasound-guided transbronchial needle aspiration (EBUS-TBNA): the needle breakage. *Ann. Thorac. Cardiovasc. Surg.* 2014;20:567–9.
6. Vial MR, ÓConnell JO, Grosu HB, Ost DE, Eapen GA, Jimenez CA. Needle fracture during endobronchial ultrasound-guided transbronchial needle aspiration. *Am. J. Respir. Crit. Care Med.* 2016;193:213–4.
7. Hanna A, Crutu A, Baldeyrou P. Needle tip fracture, an unusual EBUS-TBNA complication. *J Bronchol Intervent Pulmonol.* 2018;25:23–4.
8. Dhillon SS, Yendamuri S. Needle assembly malfunction. An unusual complication related to endobronchial ultrasound-guided transbronchial needle aspiration. *J Bronchol Intervent Pulmonol.* 2013;20:252–5.

V. Riveiro^{a,*}, A. Golpe^a, A. Casal^a, L. Valdés^{a,b}

^a *Pneumology Department, Hospital Clínico-Universitario, Santiago de Compostela, Spain*

^b *Interdisciplinary Pneumology Research Group, Santiago de Compostela Health Research Institutions (Instituto de Investigaciones Sanitarias de Santiago de Compostela/IDIS), Santiago de Compostela, Spain*

* Corresponding author.

E-mail address: vanessa.riveiro.blanco@sergas.es (V. Riveiro).

Available online 7 May 2019

<https://doi.org/10.1016/j.pulmoe.2019.04.001>
2531-0437/

© 2019 Sociedade Portuguesa de Pneumologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Treatment of advanced EGFR-mutant NSCLC patients: Sequencing matters



Dear Editor,

The availability of several treatment options in epidermal growth factor receptor mutation-positive (EGFR M+)

non-small cell lung cancer (NSCLC), poses a challenge for clinicians as to what to consider when choosing the best treatment sequence?

First, a fast and comprehensive mutation screening is required to define subpopulations benefiting from target therapies, which should include, at present time, at least EGFR, ALK, ROS1 and BRAF genes.¹

Table 1 EGFR TKIs profile.

	Mechanism of action	Advantages	Disadvantages
1G EGFR TKI	Blocks the activity of the tyrosine kinase receptor by reversibly binding the adenosine triphosphate site	Higher efficacy than chemotherapy Better toxicity profile than platinum-based chemotherapy	Unknown benefit in uncommon mutations Lack of efficacy in presence of T790M mutation Limited data on CNS activity
Erlotinib Gefitinib		Allows dose adjustments	Does not allow dose adjustments
2G EGFR TKI	ErbB family irreversible inhibitors	Higher efficacy than chemotherapy and 1G EGFR TKI Allows dose adjustments	Lack of efficacy in presence of T790M mutation
Afatinib		Higher efficacy than 1G EGFR TKI Proven efficacy even with dose reductions Approved for uncommon mutations Low hepatic toxicity/Less drug-to-drug interactions Activity in CNS BM	Requires dose management due to the toxicity profile
Dacomitinib			Unknown activity in the SNC and in uncommon mutations
3G EGFR TKI Osimertinib	Developed to specifically and irreversibly target T790M mutation Lower activity against wild-type EGFR	Efficacy in presence of T790M mutation CNS activity Less toxicity/Fewer severe AEs	Heterogeneous resistant mechanisms OS not yet known

1/2/3G, first/second/third generation; BM, brain metastases; CNS, central nervous system; EGFR, epidermal growth factor receptor; OS, overall survival; RR, response rate; T790M, substitution of a threonine (T) with a methionine (M) at position 790 of exon 20; TKI, tyrosine kinase inhibitors; TTF, time to treatment failure.

Choosing the first-line treatment should take into account several clinical and drug-related factors, such as pharmacological profile, efficacy, toxicity and progression mechanisms. Also, the first-line option will influence the treatment sequence.

There are currently several first-line treatment options available for EGFR M+ NSCLC patients, known as first (1G), second (2G) and third-generation (3G) EGFR tyrosine kinase inhibitors (TKIs) (Table 1).

Several phase III trials demonstrated superior efficacy of erlotinib (OPTIMAL, EURTAC, ENSURE), gefitinib (First-SIGNAL, IPASS, WJTOG3405, NEJ002) and afatinib (LUX-Lung 3, LUX-Lung 6) versus standard platinum-based chemotherapy.² Toxicity profile and quality of life were also superior.³

Two head-to-head trials (LUX-Lung 7 and ARCHER 1050) compared the efficacy and safety of 1G and 2G TKIs. In the LUX-Lung 7, afatinib showed superior efficacy over gefitinib considering progression free survival (PFS), time-to-treatment failure (TTF) and the objective response rate

(ORR). Also, a positive, but not statistically significant, trend was observed on overall survival (OS), favouring afatinib.^{4,5}

The phase III ARCHER 1050 trial compared dacomitinib to gefitinib in the first-line setting, excluding patients with brain metastases (BM).⁶ Dacomitinib showed benefit on PFS and OS; however, no differences were found in ORR between groups.^{6,7}

More recently, the FLAURA trial compared 3G osimertinib and 1G EGFR gefitinib and erlotinib.⁸ Osimertinib, in the first-line setting, showed better PFS, longer median duration of response and higher, although not statistically significant, ORR.⁸ At data cut-off, OS data was immature. Until now, there are no comparative data with 2G TKIs.⁸

Most of the evidence about EGFR rare mutations, which represent 10–15% of patients,⁹ comes from afatinib studies, where afatinib showed efficacy in patients with advanced-stage lung adenocarcinomas harbouring non-classical mutations, specifically on S768I, L861Q and G719X mutations.¹⁰

Another important clinical factor for first-line treatment decision is the central nervous system (CNS) activity and the presence of BM. Data on the direct intracranial activity of 1G TKIs is limited.⁹ Regarding 2G, afatinib demonstrated efficacy in patients with asymptomatic BM and delayed CNS progression⁹; no data is available on dacomitinib.⁹ Osimertinib demonstrated activity in patients with BM at standard therapeutic doses, and a reduction in CNS progression was also demonstrated.⁹

EGFR TKIs are associated with a class-related adverse event (AE) profile due to wild-type EGFR inhibition. Although afatinib is associated with a higher percentage of severe AEs, it is well demonstrated that the availability of several doses allows dose adjustments that do not compromise drug efficacy; therefore, the discontinuation rate due to AE was similar between afatinib and gefitinib on the LUX-Lung 7 study.^{2,4,5,8} Additionally, afatinib undergoes minimal hepatic metabolism and is not a substrate for liver primarily via cytochrome P450 (CYP)-dependent enzymes, having, therefore, a lower probability for drug interactions.⁹

Despite EGFR TKIs proven efficacy, inexorable disease progression still occurs. For patients with oligometastatic progression, the treatment strategy can be to keep the EGFR TKI, and local radiotherapy should be considered; for systemic progression, treatment should be changed.³

The most common mechanism of resistance related to 1G and 2G TKIs is the acquisition of the gatekeeper mutation in EGFR T790M, which occurs in 50–70% of cases.⁹

At the time of disease progression, EGFR T790M mutation can be detected through liquid biopsy or tissue rebiopsy.¹¹ If the T790M is not present in the liquid biopsy, a tissue rebiopsy should be performed.¹¹

Since rebiopsy may present some challenges (patients may refuse or not be eligible, progression may occur in an inaccessible location), the use of a highly sensitive liquid biopsy technique, which allows higher percentage of detection and a high agreement rate with the tissue biopsy, may maximise the detection of patients with T790M mutation and, therefore, the number of patients benefiting from osimertinib as a second-line treatment.¹²

Although a high sensitivity technique for the detection of the T790M mutation is not yet accessible in all hospitals, for hospitals with access to this technique, the choice of first-line therapeutic option may rely on therapeutic sequencing.

Resistance mechanisms to osimertinib present great variability and appear to be heterogeneous, with lack of specific therapeutic options after its use.^{2,9}

Therapy sequencing is a critical point. If the first option is a 1G or 2G TKI, in patients with clinically relevant progression and confirmed T790M+, osimertinib is the second-line standard of care.¹¹ Updated OS data from LUX-Lung 7 shows that sequential therapy with afatinib followed by a 3G EGFR TKI, namely osimertinib, was effective, with a 3-year OS rate up to 90%.^{5,13}

For patients T790M-negative, after progression following 1G or 2G TKI, therapy sequencing is, for now, not applicable, and chemotherapy is the current standard treatment.²

If osimertinib is used in first-line treatment, we can guarantee access for patients that might not be able to use it in 2nd line after progression with a different mechanism than T790M, or patients without the opportunity of a 2nd line.

We are still waiting for mature data on OS and PFS2 after first-line treatment with osimertinib.⁸

Unfortunately, to date, after progression, the standard of care for these patients is only platinum-based chemotherapy. There are many ongoing trials assessing agents that target specific mutations or combinations.^{2,9,11}

With the recent approval from the European Medicines Agency of the combination chemotherapy + atezolizumab + bevacizumab, a new door has opened.

There is no single therapeutic solution for the first-line treatment of NSCLC patients with EGFR M+: depending on the patient's profile and on the access to the liquid biopsy technique that allows the detection of the T790M resistance mutation, the first-line therapeutic option will vary between sequencing approach or the use of a 3G EGFR TKI in 1st line. Optimised EGFR TKI sequencing might be the most critical determinant of OS in patients with activating EGFR mutations. Data on OS from clinical trials and real-life cohorts will help us to understand the best sequence for each individual patient.

Conflicts of interest

Fernanda Estevinho declares Having received speaking fees from AstraZeneca, Bristol-Myers Squibb, Pfizer, Pierre Fabre and Roche. Participating in advisory boards of Roche and Boehringer Ingelheim.

Margarida Felizardo declares Having received speaking fees from AstraZeneca, Merck Sharp & Dohme and Roche. Participating in advisory boards of Boehringer Ingelheim.

Gabriela Fernandes declares Having received speaking fees from AstraZeneca, Boehringer Ingelheim, Merck Sharp & Dohme and Novartis. Participating in advisory boards of Boehringer Ingelheim.

Ana Figueiredo declares Having received speaking fees from Boehringer Ingelheim, Bristol-Myers Squibb, Merck Sharp & Dohme, Bristol, Novartis, Pfizer, Pierre Fabre and Roche. Participating in advisory boards of Boehringer Ingelheim and Merck Sharp & Dohme.

José Albino Lopes declares Participating in advisory boards of Boehringer Ingelheim.

Acknowledgments

Funding for this paper was provided by Boehringer Ingelheim. Funding was used to access all necessary scientific bibliography and cover meeting expenses. Boehringer Ingelheim had no role in the writing of the paper and in the decision to submit the paper for publication.

References

1. Lindeman NI, Cagle PT, Aisner DL, Arcila ME, Beasley MB, Bernicker EH, et al. Updated molecular testing guideline for the selection of lung cancer patients for treatment with targeted tyrosine kinase inhibitors: guideline from the College of American Pathologists, the International Association for the Study of Lung Cancer, and the Association for Molecular Pathology. *Arch Pathol Lab Med.* 2018;142:321–46. <http://dx.doi.org/10.5858/arpa.2017-0388-CP>.

2. Hirsh V, Turning EGFR. mutation-positive non-small-cell lung cancer into a chronic disease: optimal sequential therapy with EGFR tyrosine kinase inhibitors. *Ther Adv Med Oncol.* 2018;10, <http://dx.doi.org/10.1177/1758834017753338>.
 3. Mazza V, Cappuzzo F, Treating EGFR. mutation resistance in non-small cell lung cancer – role of osimertinib. *Appl Clin Genet.* 2017;10:49–56, <http://dx.doi.org/10.2147/TACG.S103471>.
 4. Park K, Tan EH, O'Byrne K, Zhang L, Boyer M, Mok T, et al. Afatinib versus gefitinib as first-line treatment of patients with EGFR mutation-positive non-small-cell lung cancer (LUX-Lung 7): a phase 2B, open-label, randomised controlled trial. *Lancet Oncol.* 2016;17:577–89, [http://dx.doi.org/10.1016/S1470-2045\(16\)30033-X](http://dx.doi.org/10.1016/S1470-2045(16)30033-X).
 5. Paz-Ares L, Tan EH, O'Byrne K, Zhang L, Hirsh V, Boyer M, et al. Afatinib versus gefitinib in patients with EGFR mutation-positive advanced non-small-cell lung cancer: overall survival data from the phase IIb LUX-Lung 7 trial. *Ann Oncol.* 2017;28:270–7, <http://dx.doi.org/10.1093/annonc/mdw611>.
 6. Wu YL, Cheng Y, Zhou X, Lee KH, Nakagawa K, Niho S, et al. Dacomitinib versus gefitinib as first-line treatment for patients with EGFR-mutation-positive non-small-cell lung cancer (ARCHER 1050): a randomised, open-label, phase 3 trial. *Lancet Oncol.* 2017;18(11):1454–66, [http://dx.doi.org/10.1016/S1470-2045\(17\)30608-3](http://dx.doi.org/10.1016/S1470-2045(17)30608-3).
 7. Mok TS, Cheng Y, Zhou X, Lee KH, Nakagawa K, Niho S, et al. Improvement in overall survival in a randomized study that compared dacomitinib with gefitinib in patients with advanced non-small-cell lung cancer and EGFR-activating mutations. *J Clin Oncol.* 2018;36:2244–50, <http://dx.doi.org/10.1200/JCO.7994201878>.
 8. Soria JC, Ohe Y, Vansteenkiste J, Reungwetwattana T, Chewaskulyong B, Lee KH, et al. Osimertinib in untreated EGFR-mutated advanced non-small-cell lung cancer. *N Engl J Med.* 2018;378:113–25, <http://dx.doi.org/10.1056/NEJMoa1713137>.
 9. Girard N. Optimizing outcomes in EGFR mutation-positive NSCLC: which tyrosine kinase inhibitor and when? *Future Oncol.* 2018, <http://dx.doi.org/10.2217/fon-2017-0636>.
 10. Giotrif, summary of product characteristics; 2018. http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/002280/WC500152392.pdf [accessed 05.09.18].
 11. Novello S, Barlesi F, Califano R, Cufer T, Ekman S, Levra MG, et al. ESMO guidelines committee metastatic non-small-cell lung cancer: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol.* 2016;27:v1–27.
 12. Hochmair MJ, Buder A, Schwab S, Burghuber OC, Prosch H, Hilbe W, et al. Liquid-biopsy-based identification of EGFR T790M mutation-mediated resistance to afatinib treatment in patients with advanced EGFR mutation-positive NSCLC, and subsequent response to osimertinib. *Target Oncol.* 2018;12, <http://dx.doi.org/10.1007/s11523-018-z.130612>.
 13. Corral J, Park K, Yang JCH, Mok T, Tan EH, O'Byrne K, et al. Afatinib (A) vs gefitinib (G) in patients with EGFR mutation-positive (EGFRm+) NSCLC: updated OS data from the phase IIb trial LUX-Lung 7 (LL7). *Ann Oncol.* 2017;28:ii28–51, [10./annonc/mdx091.0131093](http://dx.doi.org/10.1093/annonc/mdx091.0131093).
- F. Estevinho^{a,*}, M. Felizardo^b, G. Fernandes^c,
A. Figueiredo^d, J.A. Lopes^e
- ^a *Oncology Department, Unidade Local de Saúde de Matosinhos, EPE - Hospital Pedro Hispano, Matosinhos, Portugal*
^b *Pulmonology Department, Hospital Beatriz Ângelo, Loures, Portugal*
^c *Pulmonology Department, Centro Hospitalar e Universitário de São João, EPE, Porto, Portugal*
^d *Pulmonology Department; Centro Hospitalar e Universitário de Coimbra, EPE - Hospital Geral, Coimbra, Portugal*
^e *Pulmonology Department, Unidade Local de Saúde do Alto Minho, EPE - Hospital de Santa Luzia, Viana do Castelo, Portugal*
- *Corresponding author at: Oncology Department, Unidade Local de Saúde de Matosinhos - Hospital Pedro Hispano, Matosinhos, Rua Dr. Eduardo Torres, 4464-513, Senhora da Hora, Portugal
E-mail address: fernandaestevinho@gmail.com (F. Estevinho).
 Available online 7 June 2019
- <https://doi.org/10.1016/j.pulmoe.2019.04.0052531-0437/>
 © 2019 Sociedade Portuguesa de Pneumologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).