



PULMONOLOGY

www.journalpulmonology.org



THEMATIC SERIES

Assisted mobilisation in critical patients with COVID-19

M. Polastri^{a,b,#}, F. Daniele^b, F. Tagariello^b

^a Department of Continuity of Care and Disability, Physical Medicine and Rehabilitation, St Orsola University Hospital, Bologna, Italy

^b Department of Clinical, Integrated and Experimental Medicine (DIMES), Respiratory and Critical Care Unit, Alma Mater Studiorum University of Bologna, Bologna, Italy

Received 5 January 2021; accepted 20 January 2021

KEYWORDS

COVID-19;
Exercise;
Mobilisation;
Respiratory intensive care unit;
Respiratory failure;
Rehabilitation;
Safety

Abstract The therapeutic value of early physiotherapeutic treatment in critical respiratory settings has already been clearly outlined in the last fifteen years by several authors. However, there is still a controversial perception of mobilisation by healthcare professions.

In-bed cycling has attracted increasing attention having been demonstrated as a feasible and safe intervention in critical settings. Patients with respiratory diseases are typically prone to fatigue and exertional dyspnoea, as we observe in COVID-19 pandemic; in fact, these patients manifest respiratory and motor damage that can even be associated with cognitive and mental limitations. COVID-19 is at risk of becoming a chronic disease if the clinical sequelae such as pulmonary fibrosis are confirmed as permanent outcomes by further analysis, particularly in those cases with overlapping pre-existent pulmonary alterations.

In the present article, we propose a practical analysis of the effects of in-bed cycling, and further discuss its potential advantages if used in critical patients with COVID-19 in intensive care settings.

© 2021 Sociedade Portuguesa de Pneumologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Clinical case

A 75-years-old man with a body mass index of 26.1 kg/m² came to our attention having several comorbidities (hypertension, dyslipidemia, bronchial asthma, ulcerative colitis, critical stenosis of both the right coronary artery and the anterior intraventricular artery treated with angioplasty and stenting) and a 40 pack/year smoking history. Before hospitalisation, his daily autonomy was limited by the

Series: How I manage the case. Series Editor: Stefano Nava.
* Corresponding author at: Department of Continuity of Care and Disability, Physical Medicine and Rehabilitation, St Orsola University Hospital, Bologna, Italy.
E-mail address: massimiliano.polastri@aosp.bo.it (M. Polastri).

<https://doi.org/10.1016/j.pulmoe.2021.01.004>

2531-0437/© 2021 Sociedade Portuguesa de Pneumologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article in press as: M. Polastri, F. Daniele and F. Tagariello, Pulmonology, <https://doi.org/10.1016/j.pulmoe.2021.01.004>



Fig. 1 Multiple ground-glass areas (red line) with peri-bronchial distribution.

reduced walking capacity resulting from an accidental fall at home requiring bed rest because of limb pain and soreness. The patient went to the emergency room given the persistence of dyspnoea, cough and fever, although therapy with antibiotics and steroids had been administered at home. Hemogasanalysis showed moderate gas exchange impairment ($\text{PaO}_2/\text{FiO}_2 = 238$ mmHg and A-a O_2 gradient = 50 mmHg). The chest high-resolution computed tomography demonstrated the presence of multiple large areas of peri-bronchial ground glass compatible with bilateral interstitial pneumonia (Fig. 1). Blood tests demonstrated elevation of phlogosis indices and associated lymphopenia: Protein-C-Reactive (8.2 mg/dL), IL-6 (160.9 pg/mL), High Ferritinemia (1098 ng/mL), Lactate dehydrogenase (461 U/L), Negative Procalcitonin (0.1 ng/mL). Given the suggestive picture, a nasal/oropharyngeal swab for SARS-Cov-2 was performed and resulted positive. This case describes a patient with severe respiratory failure in bilateral interstitial pneumonia associated with critical deterioration of the physical function because of prolonged bed rest.

The patient was initially admitted to a general ward; gas-exchanges gradually deteriorated ($\text{PaO}_2/\text{FiO}_2 = 115$ mmHg). He needed oxygen therapy with Ventimask and reservoir to maintain sufficient oxygenation. Given the severe deterioration, high-dose steroid therapy was also initiated. On hospital day 8, the patient was transferred to the Respiratory Intensive Care Unit (St Orsola University Hospital, Bologna, Italy) starting high flow oxygen therapy (flow 50L/min – FiO_2 43%) because of the persistence of severe gas exchange impairment. Given the respiratory failure, medical treatment was completed with Ruxolitinib, a selective Janus Associated Kinases inhibitor usually used for haematological diseases^{1,2}. There was a gradual improvement in gas-exchanges ($\text{PaO}_2/\text{FiO}_2 = 152, 236, 278$ mmHg on hospital days 18, 20, and 21, respectively), allowing suspension of oxygen therapy. Severe sarcopenia contributed to disability and reduced ability to cope with the stress of an illness, in the present case³.

Physiotherapeutic treatment

Physical evaluation revealed difficulties in getting into the sitting position on the edge of the bed autonomously; assistance was necessary to complete the postural passage from a supine to a sitting position. Marked hypotrophy of lower

extremities was also present; hip flexion, knee extension, and ankle dorsiflexion were 3/5 as assessed by the Medical Research Council Muscle Scale⁴. A trial for a passage from a sitting to a standing position revealed the patient could not finalise the movement even with assistance of two operators. The patient started a physiotherapeutic programme consisting of in-bed motor exercises and specific postural transfer training, aiming to achieve sitting position as soon as possible.

Considering the clinical frame and the concomitant positivity to SARS-CoV-2, it was decided to use in-bed electrical cycle ergometry (MOTOmed letto2, RECK-Technik GmbH & Co. KG, Betzenweiler, Germany) to facilitate and enhance lower limb function. The device is designed to provide training for both upper and lower extremities in a passive, motor-assisted, and active mode directly from the hospital bed (Fig. 2). The patient had one daily session lasting ~30 min, depending on clinical conditions. The exercise was supervised and was to be interrupted in the case of increased heart rate (>20 bpm from the baseline) or subjective fatigue/dyspnoea higher than moderate –scoring 3– in the modified Borg Scale⁵. Active movements of upper and lower limbs were also proposed, and the patient was asked to repeat exercises autonomously during the day while in a sitting position. The patient kept to all therapeutic sessions, and no adverse events related to treatment were observed. The patient was discharged to a low-intensity setting of care on hospital day 26, being able to maintain a sitting position and having increased lower limb mobility although not yet able to reach a standing position.

Rehabilitation in critical settings: from the origins to 2020

The positive effects of early mobilisation were first described in the last century when Dr Powers, in his report, illustrated that benefits occurred in a series of 100 patients undergoing different types of surgery⁶. In that study, the treatment consisted of encouraging patients to sit out of bed and to walk already on the first postoperative day, assuming such activities would bring significant advantages in enhanced recovery and restoration of daily autonomy. The study results did indeed demonstrate that all those patients treated with early mobilisation had reduced number of days in bed, length of stay and convalescence duration in weeks.

Enhanced recovery started to be discussed as early as in the 1940s⁶; nevertheless, there were no defined and specific programmes available for implementation in critical respiratory settings until the first studies were published in the mid-1990s^{7,8}. If early postoperative rehabilitation is a valuable intervention, from that study it can be gathered that a structured rehabilitation programme is feasible, safe, and effective to counteract the physical impairment and to restore functional ability in those patients who are prone to deconditioning related to respiratory failure⁸. It should be highlighted that the main differences between postoperative rehabilitation and pulmonary rehabilitation are primarily associated with the presence of the underlying respiratory disease, which poses several challenges in terms of medical and rehabilitative issues and hospital readmis-




Fig. 2 Exercise with the patient lying in a supine position (A); exercise in a sitting position with the cycle ergometer positioned laterally to the bed (B), distancing between patient and physiotherapist is possible during the exercise session (A and B).

sion rates⁹. At the same time, it should not be forgotten that pulmonary rehabilitation is a multidisciplinary intervention involving several medical professionals to manage all the clinical aspects connected to respiratory-related physical and psychological limitations^{10,11}.

The therapeutic value of early physiotherapeutic treatment in critical respiratory settings has already been clearly outlined in the last fifteen years by several authors^{12–18}; despite this, there is still a controversial perception about mobilisation by healthcare professions¹⁹.

Searching literature

We used in-bed cycling; to date, no studies have researched the physiological effects of this type of assisted mobilisation in patients with COVID-19. However, in-bed cycling has attracted increasing attention, having been demonstrated as a feasible and safe intervention in critical settings²⁰. To retrieve evidence regarding the use of in-bed cycling we searched PubMed and Scopus databases using the following key words “in-bed cycling”, “critical patients”, and “motomed letto2”. Databases were searched from their inception until December 2020; we considered eligible only those citations written in English, Spanish, Italian or French. No limits were imposed for age, gender, and article types. We also searched the references of retrieved papers, and we made a further analysis of electronic search engines (i.e., Google) retrieving additional information.

Effects of in-bed cycling

Patients with respiratory diseases are typically prone to fatigue and exertional dyspnoea as we observe in COVID-19; in fact, these patients manifest respiratory and motor damages that can even be associated with cognitive and mental limitations²¹. COVID-19 is at risk of becoming a chronic disease if the clinical sequelae such as pulmonary fibrosis, are confirmed as permanent outcomes by further analysis^{22–27} particularly in those cases overlapping pre-existent pulmonary alterations²⁸. Patients forced to prolonged bed rest

or presenting with mobility restrictions, may simply develop muscle dysfunction or a more severe condition, namely intensive care unit acquired weakness, characterised by generalised muscle weakness affecting peripheral and respiratory muscles^{29,30}. Loss of muscle mass is negatively correlated with 2–3 weeks of immobilisation or intensive care unit stay³¹. That said, in all those conditions exposing patients to prolonged or forced restricted mobility or –even worse – immobility, primary rehabilitative goals are preserving, and hopefully, restoring a certain degree of functional capacity. COVID-19 has posed several barriers to implementing traditional physiotherapeutic activities because of the need to contain infection risk among healthcare professionals. Particularly at the beginning of the pandemic, with reduced availability of personal protective equipment, proximity with patients was conditioned by considering the risks and benefits of specific procedures, including mobilisation. A specific research line investigating characteristics and indications of personal protective equipment has contributed to improving personal protection among healthcare workers over time^{32–34}.

Can in-bed cycling in patients with COVID-19 be compared to treatments traditionally adopted in intensive care settings?

In a study by Ringdal et al., conducted to investigate physiological effects on in-bed cycling, participants – who were patients hospitalised in an intensive setting – did not show significant and/or alarming changes in physiological parameters while exercising. Heart rate, mean arterial pressure, respiratory rate, and peripheral oxygen saturation were confined to a small range of variation, confirming in-bed cycling to be safe and feasible³⁵. In the same way, participants did not manifest cardiac rhythm changes such as atrial fibrillation or ventricular tachycardia; in that study, the in-bed cycle was a suitable means for enhancing recovery and promoting motor activities in those patients who were not able to perform postural transfers³⁵. Additional physiological effects were related to interrupting sedation while cycling, allowing patients to be aware of their physical activity. Other aspects of in-bed cycling executed in critical settings are related to the patient’s perception of a feel-

Table 1 Effects of in-bed cycling.

Authors, ref. and year	N. of patients, [setting]	Main outcomes
Woo et al. ³⁹ 2018	10, [ICU]	Increase of the rectus femoris muscle cross-sectional area, and augmented circumference of the thigh
Fossat et al. ³⁸ 2018	158, [medical-surgical ICU]	The authors did not find significant differences regarding muscle strength between those patients who received in-bed cycling and electrical nerve stimulation and controls, as assessed by the MRC scale. Nor were differences between groups found for the following variables: 1) functional autonomy at discharge as assessed by the ICU Mobility Scale and the Katz Index of Independence in ADL, 2) number of ventilator-free days, 3) thickness of the rectus femoris
Machado et al. ⁴⁰ 2017	22, [general-cardiac ICU]	Increase of the peripheral muscle strength as assessed by the MRC scale
Rocca et al. ⁴¹ 2016	9, [neurological ICU]	Increase in catecholamine production in patients treated with in-bed cycling
Burtin et al. ³⁶ 2009	31, [medical-surgical ICU]	At ICU discharge the following variables were higher in patients who underwent in-bed cycling: the quadriceps force (N.kg), the walked distance (6MWT), the SF-36 PF score

Legend: ICU, intensive care unit; MRC, Medical Research Council; ADL, activities of daily living; N.kg, Newton-kilogram; 6MWT, 6-min walking test; SF-36 PF, Short Form-36 physical functioning.

ing of control, safety, and hope of recovery during a critical illness³⁵.

Several studies have shown that in-bed cycling enhances physical function, helping to prevent muscle atrophy, and it is a feasible intervention to get patients accustomed to exercise and mobility (Table 1).

In addition, in-bed cycling has been found to improve consciousness and perception of personal mobility potential, emphasising another important aspect related to the hospital stay in critical settings where patients are often subjected to sedation and mechanical ventilation^{21,41}.

Technical aspects of assisted mobilisation with in-bed cycle ergometers

In-bed cycling with the ergometer MOTomed letto2 (RECK-Technik GmbH & Co. KG, Betzenweiler, Germany) – the most widely described in the literature – ^{21,36–43} can be adopted for exercising both upper and lower extremities either in a supine or in a sitting position (Fig. 2). The cycle ergometer device can be placed either laterally or at the end of the bed and firmly anchored to the floor thanks to a blocking mechanism – preventing unwanted movements of the machine during the exercise session. The hands and legs grips/pedals are adjustable and facilitate patient positioning even in those subjects with reduced autonomous movement. The lower limbs are sustained by supports, and the feet are secured by velcro strips; hand pockets are also available for those unable to grip by themselves. The possibility of choosing a preferred exercise position allows this system to be used even in reduced spaces. The machine is equipped with wheels and can be easily moved within the ward. The cycle ergometer detects the patient's active contribution and adjusts passive or active cycling accordingly. A digital controller mounted on a movable arm allows maintaining a distance from the patient – while exercising – of approxi-

mately two meters. This feature is handy when working in COVID-19 settings because it guarantees a safe distance for the operator (Fig. 2). Once the patient has been positioned for executing the preferred exercise, they can be supervised without the need to be in strict contact; meanwhile, the physiotherapist can treat other patients nearby. When the use of in-bed cycling is optimised to the workload, one can have more slots available. Depending on the patient's clinical condition, when the cycle ergometer is placed laterally, the subject is facilitated to exercise the upper extremities and control the trunk position. Such modality is expected to enhance mobility and proprioceptive perception of the body posture requiring a certain degree of collaboration by the patient. In a way, when exercise is proposed in a sitting position on the edge of the bed, there are several advantages such as 1) improved active participation of the subject, 2) increased work of the antigravity muscles (i.e., paravertebral and abdominal groups) which can be particularly useful during weaning from bed rest related to other respiratory conditions such as either invasive and noninvasive mechanical ventilation, 3) augmented aerobic effort because of the possibility of increasing the speed, duration, intensity and type of exercise – passing from a passive/assisted to an active modality during the same session.

RT300-SUPINE (<https://www.cyclonemobility.com/product/rt300-supine-2>; Cyclone Mobility, Widnes, UK) is another cycle ergometer that incorporates functional electrical stimulation using adhesive pads. The device is mounted on an expandable wheeled table and provides arm and leg cycling; adjustable housings are present for optimising arm and leg exercise, while a dedicated system controls the legs' lateral movements during cycling. The use of the RT300-SUPINE has been described in the study by Kho et al.⁴⁴.

The THERA-Trainer Bemo (<https://thera-trainer.com/en/thera-trainer-products/cycling/thera-trainer-bemo>; medica Medizintechnik GmbH, Hochdorf, Germany) is

Table 2 In-bed cycling: training protocols.

Author, ref. and year	Frequency (days/week)	Duration (min)	Intensity	Criteria for interrupting exercise
Kho et al. ⁴⁴ 2019	–	30	–	Mean arterial pressure <60 or >110 mmHg, heart rate <40 or >140 bpm, oxygen saturation <88%
Woo et al. ³⁹ 2018	–	20	Variable	Downgrading resistance if 10% change in heart rate or blood pressure and oxygen desaturation less than 95%
Fossat et al. ³⁸ 2018	7/7	15	Passive/active	Mean arterial pressure <65 mmHg, systolic arterial pressure >180 mmHg, heart rate >135 or <45 bpm, acute arrhythmia, need of vasopressor therapy, suspected or confirmed acute myocardial ischemia, oxygen saturation <90%, any other event or condition that requires stopping the exercise
Machado et al. ⁴⁰ 2017	5/7	20	20 cycles/min	Mean arterial pressure <60 or >125 mmHg; peripheral oxygen saturation <88%; heart rate >30 or <40 bpm
Nickels et al. ⁴² 2017	6/7	20	20 cycles/min	Increase in ventilatory support persisting >5 min. post-exercise, oxygen desaturation <88% for more than 1 min, increase in systolic blood pressure >180 mmHg for more than 2 min, increase in heart rate >140 bpm for more than 2 min, decrease in mean arterial blood pressure <60 mmHg, drop in heart rate <50 bpm for more than 2 min
Kimawi et al. ⁴³ 2017	7/7	35	Progressive	At discretion of the physiotherapist considering cardiovascular and respiratory events, and catheters dislodgment
Rocca et al. ⁴¹ 2016	5/7	30	–	–
Burtin et al. ³⁶ 2009	5/7	20	Variable (20 cycles/min in sedated patients)	Heart rate decrease >20%, systolic blood pressure >180 mmHg, decrease >20% of systolic or diastolic blood pressure, peripheral oxygen saturation <90%

another device for passive, assisted and active movements of upper and lower extremities. It is a wheeled structure that allows patients to exercise either in a supine or in a sitting position, and also provides visual feedback during the session. The device can be adjusted using the control and display unit; it includes emergency-stop functions and a system detecting spasticity.

Treatment's details of in-bed cycling

Considering the evidence available, we gathered some common treatment schemes regarding duration, type, time, and intensity of exercise using in-bed cycling, as shown in [Table 2](#). From these studies, it was possible to decide which criteria should be used for interrupting training; standard hemodynamic parameters such as heart rate, arterial pressure, and oxygen saturation should be used to guarantee safety during in-bed cycling.

In one study by Machado et al., the authors even provided the physiotherapist-to-patient ratio 1:8 returning an interesting consideration on physiotherapists' workload⁴⁰.

Synthesising data from [Table 2](#), exercise frequency can range between 5 to 7 days a week, duration between 5 to 35 min, with variable intensity – passive, assisted, active – depending on patients' clinical conditions.

Is in-bed cycling feasible and safe?

Exercise with an in-bed cycle ergometer has been demonstrated to be feasible and safe, as outlined in several types of research^{20,37,43,44}. In a recent study, overall comments from patients, clinicians, and family members were positive when recalling cycling sessions; all were concordant, affirming that in-bed cycling contributed to physical recovery²⁰.

Among the effects of in-bed cycling, the sympathetic system stimulation has been found by Rocca et al. in patients with severe neurological injury⁴¹; such a stimulation seems to be recognised as a mechanism leading to increased production of catecholamines. The authors concluded that in-bed cycling, in patients with subarachnoid haemorrhage, should be used with caution because of the potential risk of vasospasm due to elevated catecholamine levels. However, they did not find contraindications to its use to prevent polyneuromyopathy in critical illness⁴¹. In intensive care settings, in-bed cycling should be supervised to guarantee safety during exercise at all times ([Table 3](#)).

Further considerations

In-bed cycle ergometry seems to be a promising intervention in different types of patients including intensive care

Table 3 In-bed cycling: adverse events reported in the literature.

Author, ref. and year	N. of patients, [setting]	Adverse events
Nickels et al. ²⁰ 2020	36 patients, [mixed ICU]. Participants were enrolled if expected to be mechanically ventilated >48 h and to remain in ICU >48 h	Two minor events occurred – in the same patient – during 276 cycling sessions consisting of oxygen desaturation and increased respiratory rate
Kho et al. ⁴⁴ 2019	34 patients, [ICU]. Treatment was implemented within the first 4 days of MV and 7 days of ICU admission	Patients cycled with a femoral catheter in situ, none of the subjects experienced unplanned extubation or bleeding at the catheter site. 1 out of 34 participants experienced supraventricular tachycardia requiring exercise interruption
Fossat et al. ³⁸ 2018	158 patients, [medical-surgical ICU]. Patients were enrolled if expected to remain in ICU >48 h, and with a BI >55 within the 15 days before admission	1 unplanned extubation requiring immediate reintubation
Kimawi et al. ⁴³ 2017	106 patients, [medical ICU]. A percentage of them was under mechanical ventilation, continuous renal replacement therapy, and had femoral catheters in situ during cycling	Twelve abnormalities occurred during 260 cycling sessions; those events were cardiovascular or respiratory alterations, catheter removal or dislodgment, tube (artificial airway, chest, or feeding) removal. Such circumstances did not require therapy, and they did not cause additional costs or increased length of stay
Kho et al. ³⁷ 2015	181 patients, [medical ICU]. A percentage of them was under mechanical ventilation, vasopressors infusion, and renal replacement therapy having femoral arterial, dialysis, or venous catheters in situ during some sessions of cycling	One event was represented by the dislodgment before cycling of a radial arterial line already scheduled for replacement because of unstable positioning

Legend: ICU, intensive care unit; MV, mechanical ventilation; BI, Barthel Index.

settings, as outlined in several research studies. Although such a trend has been confirmed in published studies, up until now no clinical experiments have described the use of in-bed cycling in patients with COVID-19. Considering the knowledge and evidence gathered from literature, we feel there is enough to corroborate the hypothesis that in-bed cycling can be used in COVID settings with selected patients (i.e., respiratory-related motor impairment, sarcopenia, and muscle hypotrophy related to prolonged bed rest). In addition to the therapeutic benefits for patients, in-bed cycling allows healthcare professionals to distance while operating, and this is not of secondary importance in such a pandemic. Additional advantages can be found in delegating exercise supervision to different multidisciplinary team members – once instructed – freeing specific staff – such as physiotherapists – for other simultaneous interventions, particularly when human resources are lacking or scarce.

Funding

None.

Conflict of interests

The authors have no conflicts of interest to declare.

Acknowledgements

We would like to thank Cecilia Mari for revising the use of English in the present manuscript.

References

1. La Rosée F, Bremer HC, Gehrke I, Kehr A, Hochhaus A, Birndt S, et al. The Janus kinase 1/2 inhibitor ruxolitinib in COVID-19 with severe systemic hyperinflammation. *Leukemia*. 2020;34(7):1805–15.
2. Gozzetti A, Capochiani E, Bocchia M. The Janus kinase 1/2 inhibitor ruxolitinib in COVID-19. *Leukemia*. 2020;34(10):2815–6.
3. Roubenoff R. Sarcopenia and its implications for the elderly. *Eur J Clin Nutr*. 2000;54 Suppl 3:S40–7.
4. Medical Research Council. Aids to the examination of the peripheral nervous system. Available from: <https://tinyurl.com/y5tc3ovw>.
5. Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc*. 1982;14(5):377–81.
6. Powers JH. The abuse of rest as a therapeutic measure in surgery: early postoperative activity and rehabilitation. *JAMA*. 1944;125(16):1079–83.
7. Nava S, Rubini F, Zanotti E, Ambrosino N, Bruschi C, Vitacca M, et al. Survival and prediction of successful ventilator weaning in COPD patients requiring mechanical ventilation for more than 21 days. *Eur Respir J*. 1994;7(9):1645–52.

8. Nava S. Rehabilitation of patients admitted to a respiratory intensive care unit. *Arch Phys Med Rehabil.* 1998;79(7):849–54.
9. Polastri M, Pisani L, Dell'Amore A, Nava S. Revolving door respiratory patients: a rehabilitative perspective. *Monaldi Arch Chest Dis.* 2017;87(3):857.
10. Spruit MA, Singh SJ, Garvey C, ZuWallack R, Nici L, Rochester C, et al. An Official American Thoracic Society/European Respiratory Society statement: key concepts and advances in pulmonary rehabilitation. *Am J Respir Crit Care Med.* 2013;188(8):e13–64.
11. Spruit MA, Wouters EFM. Organizational aspects of pulmonary rehabilitation in chronic respiratory diseases. *Respirology.* 2019;24(9):838–43.
12. Ceriana P, Delmastro M, Rampulla C, Nava S. Demographics and clinical outcomes of patients admitted to a respiratory intensive care unit located in a rehabilitation center. *Respir Care.* 2003;48(7):670–6.
13. Clini E, Ambrosino N. Early physiotherapy in the respiratory intensive care unit. *Respir Med.* 2005;99(9):1096–104.
14. Bailey P, Thomsen GE, Spuhler VJ, Blair R, Jewkes J, Bezdjian L, et al. Early activity is feasible and safe in respiratory failure patients. *Crit Care Med.* 2007;35(1):139–45.
15. Morris PE, Griffin L, Berry M, Thompson C, Hite RD, Winkelman C, et al. Receiving early mobility during an intensive care unit admission is a predictor of improved outcomes in acute respiratory failure. *Am J Med Sci.* 2011;341(5):373–7.
16. Gosselink R, Bott J, Johnson M, Dean E, Nava S, Norrenberg M, et al. Physiotherapy for adult patients with critical illness: recommendations of the European Respiratory Society and European Society of Intensive Care Medicine Task Force on physiotherapy for critically ill patients. *Intensive Care Med.* 2008;34(7):1188–99.
17. Ceriana P, Ambrosino N. Rehabilitation in the intensive care unit. In: Donner CF, Ambrosino N, Goldstein R, editors. *Pulmonary rehabilitation.* 2nd edition CRC Press; 2020. p. 391–8.
18. Martí JD, McWilliams D, Gimeno-Santos E. Physical therapy and rehabilitation in chronic obstructive pulmonary disease patients admitted to the intensive care unit. *Semin Respir Crit Care Med.* 2020;41(6):886–98.
19. Hermes C, Nydahl P, Blobner M, Dubb R, Filipovic S, Kaltwasser A, et al. Assessment of mobilization capacity in 10 different ICU scenarios by different professions. *PLoS One.* 2020;15(10):e0239853.
20. Nickels MR, Aitken LM, Barnett AG, Walsham J, McPhail SM. Acceptability, safety, and feasibility of in-bed cycling with critically ill patients. *Aust Crit Care.* 2020;33(3):236–43.
21. Simpson R, Robinson L. Rehabilitation after critical illness in people with COVID-19 infection. *Am J Phys Med Rehabil.* 2020;99(6):470–4.
22. Atabati E, Dehghani-Samani A, Mortazavimoghaddam SG. Association of COVID-19 and other viral infections with interstitial lung diseases, pulmonary fibrosis, and pulmonary hypertension: a narrative review. *Can J Respir Ther.* 2020;56:1–9.
23. Fang Y, Zhou J, Ding X, Ling G, Yu S. Pulmonary fibrosis in critical ill patients recovered from COVID-19 pneumonia: preliminary experience. *Am J Emerg Med.* 2020;38(10):2134–8.
24. Wigén J, Löfdahl A, Björmer L, Elowsson-Rendin L, Westergren-Thorsson G. Converging pathways in pulmonary fibrosis and Covid-19—the fibrotic link to disease severity. *Respir Med X.* 2020;2:100023.
25. Yang ZL, Chen C, Huang L, Zhou SC, Hu YN, Xia LM, et al. Fibrotic changes depicted by thin-section CT in patients with COVID-19 at the early recovery stage: preliminary experience. *Front Med (Lausanne).* 2020;7:605088.
26. Picchi G, Mari A, Ricciardi A, Carucci AC, Sinatti G, Cosimini B, et al. Three cases of COVID-19 pneumonia in female patients in Italy who had pulmonary fibrosis on follow-up lung computed tomography imaging. *Am J Case Rep.* 2020;21:e926921.
27. Tale S, Ghosh S, Meitei SP, Kolli M, Garbhapu AK, Pudi S. Post-COVID-19 pneumonia pulmonary fibrosis. *QJM.* 2020;113(11):837–8.
28. Mazzolini M, Monari M, Angeletti G, Dalpiaz G, Rocca A. Fatal pulmonary fibrosis complicating COVID-19 infection in preexistent emphysema. *Radiol Case Rep.* 2021;16(2):361–3.
29. Hermans G, Van den Bergh G. Clinical review: intensive care unit acquired weakness. *Crit Care.* 2015;19(1):274.
30. Kress JP, Hall JB. ICU acquired weakness and recovery from critical illness. *N Engl J Med.* 2014;370(17):1626–35.
31. Gruther W, Benesch T, Zorn C, Paternostro-Sluga T, Quittan M, Fialka-Moser V, et al. Muscle wasting in intensive care patients: ultrasound observation of the M. quadriceps femoris muscle layer. *J Rehabil Med.* 2008;40(3):185–9.
32. Ippolito M, Vitale F, Accurso G, Iozzo P, Gregoretti C, Giarratano A, et al. Medical masks and respirators for the protection of healthcare workers from SARS-CoV-2 and other viruses. *Pulmonology.* 2020;26(4):204–12.
33. Ferioli M, Cisternino C, Leo V, Pisani L, Palange P, Nava S. Protecting healthcare workers from SARS-CoV-2 infection: practical indications. *Eur Respir Rev.* 2020;29(155):200068.
34. Winck JC, Ambrosino N. COVID-19 pandemic and non invasive respiratory management: every Goliath needs a David. An evidence based evaluation of problems. *Pulmonology.* 2020;26(4):213–20.
35. Ringdal M, Warren Stomberg M, Egnell K, Wennberg E, Zätterman R, Rylander C. In-bed cycling in the ICU; patient safety and recollections with motivational effects. *Acta Anaesthesiol Scand.* 2018;62(5):658–65.
36. Burtin C, Clerckx B, Robbeets C, Ferdinande P, Langer D, Troosters T, et al. Early exercise in critically ill patients enhances short-term functional recovery. *Crit Care Med.* 2009;37(9):2499–505.
37. Kho ME, Martin RA, Toonstra AL, Zanni JM, Manthey EC, Nelliot A, et al. Feasibility and safety of in-bed cycling for physical rehabilitation in the intensive care unit. *J Crit Care.* 2015;30(6), 1419.e1-5.
38. Fossat G, Baudin F, Courtes L, Bobet S, Dupont A, Bretagnol A, et al. Effect of in-bed leg cycling and electrical stimulation of the quadriceps on global muscle strength in critically ill adults: a randomized clinical trial. *JAMA.* 2018;320(4):368–78.
39. Woo K, Kim J, Kim HB, Choi H, Kim K, Lee D, et al. The effect of electrical muscle stimulation and in-bed cycling on muscle strength and mass of mechanically ventilated patients: a pilot study. *Acute Crit Care.* 2018;33(1):16–22.
40. Machado ADS, Pires-Neto RC, Carvalho MTX, Soares JC, Cardoso DM, Albuquerque IM. Effects that passive cycling exercise have on muscle strength, duration of mechanical ventilation, and length of hospital stay in critically ill patients: a randomized clinical trial. *J Bras Pneumol.* 2017;43(2):134–9.
41. Rocca A, Pignat JM, Berney L, Jöhr J, Van de Ville D, Daniel RT, et al. Sympathetic activity and early mobilization in patients in intensive and intermediate care with severe brain injuries: a preliminary prospective randomized study. *BMC Neurol.* 2016;16(1):169.
42. Nickels MR, Aitken LM, Walsham J, Barnett AG, McPhail SM. Critical care cycling study (CYCLIST) trial protocol: a randomised controlled trial of usual care plus additional in-bed cycling sessions versus usual care in the critically ill. *BMJ Open.* 2017;7(10):e017393.
43. Kimawi I, Lamberjack B, Nelliot A, Toonstra AL, Zanni J, Huang M, et al. Safety and feasibility of a protocolized approach to in-bed cycling exercise in the intensive care unit: quality improvement project. *Phys Ther.* 2017;97(6):593–602.
44. Kho ME, Molloy AJ, Clarke FJ, Reid JC, Herridge MS, Karachi T, et al. Multicentre pilot randomised clinical trial of early in-bed cycle ergometry with ventilated patients. *BMJ Open Respir Res.* 2019;6(1):e000383.