



EDITORIAL

Asthma hospitalizations: A call for a national strategy to fight health inequities



Health inequities are differences in the health status or the distribution of health resources between different population groups, arising from the social conditions in which people are born, grow, live, work and age.¹ Such inequities have high social and economic costs for individuals and societies worldwide.¹ Lower socioeconomic status is among the risk factors associated with worse health outcomes and leads to a higher rate of potentially preventable hospital admissions.² This may reflect regional differences in access to healthcare (e.g., delays in access to secondary care) and primary care quality, which have been significantly associated with avoidable emergency admissions.³⁻⁵ This is especially important for chronic conditions in which proper management may avoid exacerbations,^{6,7} as is the case of asthma.⁸ There is evidence that people with asthma living in deprived areas may have impaired asthma outcomes across all stages of patient care, leading to a higher rate of hospitalizations and risk of asthma-related deaths.^{9,10} As health inequities have remained insufficiently studied in Portugal, in particular regarding access to care,¹¹ we assessed whether such inequities may occur in Portugal as well, by studying the association between the region of residence and the frequency of asthma hospitalizations.

In mainland Portugal, between 2011 and 2015, there were 9161 asthma hospitalizations in adults, according to data from the national administrative database containing all hospitalizations in public hospitals from mainland Portugal (ICD-9-CM code 493.x). Therefore, there was an average of 92.6 hospital admissions per 100,000 inhabitants and 13.5 hospital admissions per 1000 inhabitants with asthma. Mainland Portugal is composed of 18 districts (Fig. 1A). The districts with the highest rates of asthma hospital admissions per 100,000 inhabitants were Coimbra (213.2 hospitalizations per 100,000 inhabitants), Castelo Branco (160.6/100,000), Viseu (155.4/100,000) and Guarda (140.9/100,000), all of which are in the center region of Portugal and are (except for Coimbra) inland districts. Taking into account the prevalence data from the most recent Portuguese epidemiological survey carried out in 2010¹² (Fig. 1B),

the three districts with the most prevalence-adjusted asthma hospital admissions were Guarda (82.9 hospitalizations per 1000 inhabitants with asthma), Castelo Branco (32.8/1000), and Viseu (29.9/1000). All these three districts have a low urban coverage. Importantly, Coimbra, an urban district in coastal Portugal, is not among the districts with the highest rate of prevalence-adjusted hospital admissions, despite being among the districts displaying the highest crude hospitalization rate for asthma.

The inland regions of Portugal are, on average, less urbanized than its coastal regions,¹³ and Guarda, Viseu and Castelo Branco are among the districts with the least purchasing power (according to data from the National Institute of Statistics¹⁴ and defined according to its published definition¹⁵), so it is possible that such results could be partially explained by socioeconomic disadvantage. It is also known that the population in inland Portugal is, on average, older than its coastal counterpart. Guarda and Castelo Branco are in the top three districts with the highest aging index (proportion of inhabitants aged 65 years or older relative to those younger than 15 years older) in Portugal.¹⁶ Several comorbidities are associated with asthma in the elderly, which may complicate asthma management in these patients.¹⁷ Older age is also associated with lower literacy, which may hinder adherence to treatment in chronic diseases in elderly people.¹⁸ As a result, differences in age distribution between districts in mainland Portugal may partially explain the differences in hospital admissions for asthma. Additionally, according to data from the Portuguese Medical Association, districts in inland Portugal, especially Guarda and Castelo Branco, are also among those with fewer registered physicians per 100,000 inhabitants.¹⁹

Nevertheless, even when adjusting for purchasing power, aging index and density of registered physicians in Portuguese districts, these three districts still showed the highest number of prevalence-adjusted asthma hospital admissions (Fig. 1D-F; Table 1). This suggests potential regional health inequities that are not solely explained by factors such as age and purchasing power and may reflect the ongoing asthma

<https://doi.org/10.1016/j.pulmoe.2022.12.001>

2531-0437/© 2023 Sociedade Portuguesa de Pneumologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1 Data on (A) population density per district in mainland Portugal (1 – Viana do Castelo, 2 – Braga, 3 – Porto, 4 – Vila Real, 5 – Bragança, 6 – Aveiro, 7 – Viseu, 8 – Guarda, 9 – Coimbra, 10 – Castelo Branco, 11 – Leiria, 12 – Lisboa, 13 – Santarém, 14 – Portalegre, 15 – Setúbal, 16 – Évora, 17 – Beja, 18 – Faro); (B) asthma prevalence in mainland Portugal (percentage), and on asthma hospital admissions in mainland Portugal (C) per 1000 patients with asthma; (D) per 1000 patients with asthma and adjusted for a purchasing power of 100% *per capita*; (E) per 1000 patients with asthma and adjusted for a standard population with 1000 physicians per 100,000 inhabitants; (F) per 1000 patients with asthma and adjusted for a purchasing power of 100% *per capita* and a standard population with 1000 physicians per 100,000 inhabitants; and (G) per 1000 patients with asthma and adjusted for a purchasing power of 100% *per capita* and a standard population with 1000 physicians per 100,000 inhabitants and an aging index of 100.

Table 1 Prevalence of asthma and hospital admissions for asthma per district of mainland Portugal.

	Prevalence of asthma, %	Hospital admissions, <i>n</i>	Hospital admissions per 100,000 inhabitants, <i>n</i>	Hospital admissions per 1000 asthmatics, <i>n</i>			
				Unadjusted	adjusted for purchasing power	adjusted for physician density	adjusted for purchasing power and physician density
Aveiro	5.0	552	78.3	15.7	17.5	55.8	62.3
Beja	9.6	86	58.0	6.0	7.2	28.3	34.0
Braga	4.2	413	36.7	8.7	10.2	26.2	30.7
Bragança	6.8	108	82.7	12.2	15.7	45.7	58.8
Castelo Branco	4.9	302	160.6	32.8	39.6	119.8	144.5
Coimbra	8.8	888	213.2	24.2	25.3	20.7	21.6
Évora	6.1	96	59.6	9.8	10.7	28.3	30.8
Faro	6.3	536	121.1	19.2	20.0	52.4	54.5
Guarda	1.7	216	140.9	82.9	108.9	298.3	392.0
Leiria	7.9	398	85.8	10.9	12.0	53.5	58.8
Lisboa	7.8	2423	108.2	13.9	10.4	20.3	15.2
Portalegre	5.6	52	46.0	8.2	9.6	24.6	28.9
Porto	7.7	1386	77.3	10.0	9.9	16.6	16.5
Santarém	9.2	302	68.0	7.4	8.3	33.1	37.2
Setúbal	4.6	501	58.8	12.8	12.8	45.0	45.0
Viana do Castelo	8.1	158	66.1	8.2	10.3	25.1	31.4
Vila Real	7.4	277	138.7	18.7	24.1	34.3	44.2
Viseu	5.2	571	155.4	29.9	39.2	121.0	158.6

management available/provided in different regions. This suggests that inequities in access to healthcare services may hinder asthma management in these inland districts. The results of this study may be a model to be implemented in other underserved populations in Europe and globally.

Early asthma diagnosis and vigilant control are crucial to prevent asthma exacerbations and reduce the healthcare burden. Initiatives in Finland, Poland and Brazil have shown that prioritizing asthma care and placing primary care at the center of asthma care reduce both asthma morbidity and mortality.^{20,21} However, the Global Initiative for Asthma (GINA) guidelines have primarily been developed and tailored by tertiary care physicians and may be challenging to implement in primary care, particularly when physicians' density is low.²²⁻²⁴ More easy-to-implement and patient-centered care pathways for asthma, including all health care professionals, are needed. Additionally, as medication adherence in asthma tends to be poor,²⁵ tailored, integrated and transdisciplinary approaches to raise asthma literacy in disfavored regions could increase adherence to medication by patients. Moreover, digital solutions may be of use to improve asthma care, especially for regions with suboptimal access to healthcare services. Telemedicine may facilitate access to asthma clinical reviews and allow for remote patient monitoring, thus preventing exacerbations.²⁶ Emerging technologies, such as monitoring devices and mobile apps, may further improve asthma care. The potential of mobile apps for patient-centered care has been previously shown for allergic rhinitis²⁷, and there are reasons to believe such may apply to asthma care as well.²⁸ Apps may therefore provide information on asthma-related triggers, such as pollen season and pollution, and feedback from mobile apps may be used by the patient and the physician to improve asthma management.²⁹

In conclusion, our results suggest the existence of health inequities in asthma management in mainland Portugal and an urgent call for action from policy makers. This issue is surmounted as depopulation and aging are increasing in rural areas of Europe.³⁰ Complementary to the traditional Health Service approaches, in these regions, digitally-enabled, patient-centered care may contribute to reducing asthma burden and hospitalizations, as well as health inequities in this population.

Funding sources

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Acknowledgements

None.

References

1. World Health Organisation. Health inequities and their causes, <https://www.who.int/news-room/facts-in-pictures/detail/health-inequities-and-their-causes>; 2018 [accessed 30 November 2022].
2. McCartney G, Hart C, Watt G. How can socioeconomic inequalities in hospital admissions be explained? A cohort study. *BMJ Open*. 2013;3(8):e002433. <https://doi.org/10.1136/bmjopen-2012-002433>.
3. McCormick B, Hill PS, Redding S. Comparative morbidities and the share of emergencies in hospital admissions in deprived areas: a method and evidence from English administrative data. *BMJ Open*. 2018;8(8):e022573. <https://doi.org/10.1136/bmjopen-2018-022573>.
4. Parkinson B, Meacock R, Checkland K, Sutton M. How sensitive are avoidable emergency department attendances to primary care quality? Retrospective observational study. *BMJ Qual Saf*. 2021;30(11):884–92. <https://doi.org/10.1136/bmjqs-2020-011651>.
5. Prentice JC, Pizer SD. Waiting times and hospitalizations for ambulatory care sensitive conditions. *Health Serv Outcomes Res Methodol*. 2008;8(1):1–18. <https://doi.org/10.1007/s10742-007-0024-5>.
6. Crook ED, Peters M. Health disparities in chronic diseases: where the money is. *Am J Med Sci*. 2008;335(4):266–70. <https://doi.org/10.1097/maj.0b013e31816902f1>.
7. Van Loenen T, Faber MJ, Westert GP, Van den Berg MJ. The impact of primary care organization on avoidable hospital admissions for diabetes in 23 countries. *Scand J Prim Health Care*. 2016;34(1):5–12. <https://doi.org/10.3109/02813432.2015.1132883>.
8. Fleetcroft R, Noble M, Martin A, Coombes E, Ford J, Steel N. Emergency hospital admissions for asthma and access to primary care: cross-sectional analysis. *Br J Gen Pract*. 2016;66(650):e640–6. <https://doi.org/10.3399/bjgp16X686089>.
9. De La Cruz N, Hines JH, Shaw C, Appiah D. Geographic disparity in asthma hospitalizations: the role of race/ethnicity, socioeconomic status, and other factors. *Cureus*. 2021;13(11):e20015. <https://doi.org/10.7759/cureus.20015>.
10. Sallakh MAA, Rodgers SE, Lyons RA, Sheikh A, Davies GA. Socioeconomic deprivation and inequalities in asthma care in Wales. *Lancet*. 2017;390:S19. [https://doi.org/10.1016/S0140-6736\(17\)32954-9](https://doi.org/10.1016/S0140-6736(17)32954-9).
11. Campos-Matos I, Russo G, Perelman J. Connecting the dots on health inequalities – a systematic review on the social determinants of health in Portugal. *Int J Equity Health*. 2016;15(1):26. <https://doi.org/10.1186/s12939-016-0314-z>.
12. Sa-Sousa A, Morais-Almeida M, Azevedo LF, et al. Prevalence of asthma in Portugal - the Portuguese National Asthma Survey. *Clin Transl Allergy*. 2012;2(1):15. <https://doi.org/10.1186/2045-7022-2-15>.
13. Ayres-Sampaio D, Teodoro AC, Sillero N, Santos C, Fonseca J, Freitas A. An investigation of the environmental determinants of asthma hospitalizations: an applied spatial approach. *Appl Geogr*. 2014;47:10–9. <https://doi.org/10.1016/j.apgeog.2013.11.011>.
14. PORDATA – Estatísticas sobre Portugal e Europa. Poder de compra per capita, <https://www.pordata.pt/Municipios/Poder+de+compra+per+capita-118>; 2022 [accessed 30 November 2022].
15. Instituto Nacional de Estatística. Estudo sobre o Poder de Compra Concelhio: 2015, <https://www.ine.pt>; 2017 [accessed 30 November 2022].
16. PORDATA – Estatísticas sobre Portugal e Europa. Indicadores de envelhecimento. <https://www.pordata.pt/Portugal/Indicadores+de+envelhecimento-526>; 2022 [accessed 30 November 2022].
17. Battaglia S, Benfante A, Spatafora M, Scichilone N. Asthma in the elderly: a different disease? *Breathe*. 2016;12(1):18. <https://doi.org/10.1183/20734735.002816>.
18. Barrett GF, Riddell WC. Ageing and skills: the case of literacy skills. *Eur J Educ*. 2019;54(1):60–71. <https://doi.org/10.1111/ejed.12324>.
19. Ordem dos Médicos. Estatísticas Nacionais, <https://ordemdosmedicos.pt/estatisticas-nacionais/>; 2022 [accessed 30 November 2022].
20. Haahela T, Tuomisto LE, Pietinalho A, et al. A 10 year asthma programme in Finland: major change for the better. *Thorax*. 2006;61(8):663–70. <https://doi.org/10.1136/thx.2005.055699>.

21. Kupczyk M, Haahtela T, Cruz AA, Kuna P. Reduction of asthma burden is possible through National Asthma Plans. *Allergy*. 2010;65(4):415–9. <https://doi.org/10.1111/j.1398-9995.2009.02265.x>.
22. Gagné ME, Boulet LP. Implementation of asthma clinical practice guidelines in primary care: a cross-sectional study based on the Knowledge-to-Action Cycle. *J Asthma*. 2018;55(3):310–7. <https://doi.org/10.1080/02770903.2017.1323919>.
23. Global Initiative for Asthma. Global strategy for asthma management and prevention, www.ginasthma.org; 2022 [accessed 30 November 2022]
24. Price C, Agarwal G, Chan D, et al. Large care gaps in primary care management of asthma: a longitudinal practice audit. *BMJ Open*. 2019;9(1):e022506. <https://doi.org/10.1136/bmjopen-2018-022506>.
25. Engelkes M, Janssens HM, de Jongste JC, Sturkenboom MCJM, Verhamme KMC. Medication adherence and the risk of severe asthma exacerbations: a systematic review. *Eur Respir J*. 2015;45(2):396. <https://doi.org/10.1183/09031936.00075614>.
26. Chongmelaxme B, Lee S, Dhippayom T, Saokaew S, Chaiyakunapruk N, Dilokthornsakul P. The effects of telemedicine on asthma control and patients' quality of life in adults: a systematic review and meta-analysis. *J Allergy Clin Immunol Pract*. 2019;7(1):199–216.e111. <https://doi.org/10.1016/j.jaip.2018.07.015>.
27. Antó A, Sousa-Pinto B, Czarlewski W, et al. Automatic market research of mobile health apps for the self-management of allergic rhinitis. *Clin Exp Allergy*. 2022;52(10):1195–207. <https://doi.org/10.1111/cea.14135>.
28. Bousquet J, Sousa-Pinto B, Anto J, et al. Identification by cluster analysis of patients with asthma and nasal symptoms using the MASK-air® mHealth app. *Pulmonology*. 2022. <https://doi.org/10.1016/j.pulmoe.2022.10.005>. Nov 22;S2531-0437(22)00252-5.
29. Kvedariene V, Biliute G, Didziokaitė G, et al. Mobile health app for monitoring allergic rhinitis and asthma in real life in Lithuanian MASK-air users. *Clin Transl Allergy*. 2022;12(9):e12192. <https://doi.org/10.1002/ctt2.12192>.
30. Camarero L, Oliva J. Thinking in rural gap: mobility and social inequalities. *Palgrave Commun*. 2019;5(1):1–7.
- R.J. Vieira^{a,b}, B. Sousa-Pinto^{a,b,*}, A.M. Pereira^{a,c}, C.R. Cordeiro^{d,e}, C.C. Loureiro^{d,e}, F. Regateiro^{f,g,h}, L. Taborda-Barata^{i,j,k}, J. Bousquet^{l,m,n,o}, J. Fonseca^{a,b,c}
- ^a Department of Community Medicine, Information and Health Decision Sciences (MEDCIDS), Faculty of Medicine, University of Porto, Portugal
- ^b Centre for Health Technology and Services Research, Health Research Network (CINTESIS@RISE), Faculty of Medicine of the University of Porto, Portugal
- ^c Allergy Unit, CUF Porto Hospital & Institute, Portugal
- ^d Department of Pulmonology, University Hospital of Coimbra, Portugal
- ^e Faculty of Medicine, University of Coimbra, Portugal
- ^f Allergy and Clinical Immunology Unit, Centro Hospitalar Universitário de Coimbra, Coimbra, Portugal
- ^g Institute of Immunology, Faculty of Medicine, University of Coimbra, Coimbra, Portugal
- ^h Coimbra Institute for Clinical and Biomedical Research (iCBER), Faculty of Medicine, University of Coimbra, Coimbra, Portugal
- ⁱ UBIAir - Clinical & Experimental Lung Centre, University of Beira Interior, Covilhã, Portugal
- ^j CICS-Health Sciences Research Centre, University of Beira Interior, Covilhã, Portugal
- ^k Department of Immunoallergology, Cova da Beira University Hospital Centre, Covilhã, Portugal
- ^l Institute for Allergology, Charité – Universitätsmedizin Berlin, Corporate, Member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany
- ^m Fraunhofer Institute for Translational Medicine and Pharmacology ITMP, Allergology and Immunology, Berlin, Germany
- ⁿ University Hospital, Montpellier, France
- ^o ARIA, Montpellier, France
- * Corresponding author at: MEDCIDS – Department of Community Medicine, Information and Health Decision Sciences (MEDCIDS), Faculty of Medicine, University of Porto, Rua Dr. Plácido da Costa s/n, 4200-450, Porto, Portugal.
- E-mail address: bernardo@med.up.pt (B. Sousa-Pinto).
- Received 22 August 2022; Accepted 1 December 2022
Available online 10 March 2023